

A CHAIN THEOREM FOR MATROIDS

James Oxley, Charles Semple and Geoff Whittle

*Department of Mathematics and Statistics
University of Canterbury
Private Bag 4800
Christchurch, New Zealand*

Report Number: UCDMS2006/13

DECEMBER 2006

A CHAIN THEOREM FOR MATROIDS

JAMES OXLEY, CHARLES SEMPLE, AND GEOFF WHITTLE

ABSTRACT. Tutte's Wheels-and-Whirls Theorem proves that if M is a 3-connected matroid other than a wheel or a whirl, then M has a 3-connected minor N such that $|E(M)| - |E(N)| = 1$. Geelen and Whittle extended this theorem by showing that when M is sequentially 4-connected, the minor N can also be guaranteed to be sequentially 4-connected, that is, for every 3-separation (X, Y) of N , the set $E(N)$ can be obtained from X or Y by successively applying the operations of closure and coclosure. Hall proved a chain theorem for a different class of 4-connected matroids, those for which every 3-separation has at most five elements on one side. This paper proves a chain theorem for those sequentially 4-connected matroids that also obey this size condition.

1. INTRODUCTION

In dealing with matroid connectivity, one frequently wants to be able to remove a small set of elements from a matroid M to obtain a minor N that maintains the connectivity of M . Such results are referred to as *chain theorems*. Tutte [10] proved that if M is 2-connected and $e \in E(M)$, then $M \setminus e$ or M/e is 2-connected. More profoundly, when M is 3-connected, Tutte [10] proved the following result, his Wheels-and-Whirls Theorem.

Theorem 1.1. *Let M be a 3-connected matroid other than a wheel or whirl. Then M has an element e such that $M \setminus e$ or M/e is 3-connected.*

This result has proved to be such a useful tool for 3-connected matroids that it is natural to seek a corresponding result for 4-connected matroids. Since higher connectivity for matroids may be unfamiliar, we now define it. For a matroid M with ground set E and rank function r , the *connectivity function* λ_M of M is defined on all subsets X of E by $\lambda_M(X) = r(X) + r(E - X) - r(M)$. A subset X or a partition $(X, E - X)$ of E is *k-separating* if $\lambda_M(X) \leq k - 1$. A *k-separating* partition $(X, E - X)$ is a *k-separation* if $|X|, |E - X| \geq k$. A matroid having no *k-separations* for all $k < n$ is *n-connected*.

For 4-connected matroids, the hope of a chain theorem is frustrated by examples given by Rajan [9]. He showed that, for all positive integers m , there

Date: November 27, 2006.

1991 Mathematics Subject Classification. 05B35.

The first author was supported by the National Security Agency and the second and third authors were supported by the New Zealand Marsden Fund.

is a 4-connected matroid M such that M has no proper 4-connected minor N with $|E(M)| - |E(N)| \leq m$. Rajan also supplied corresponding examples for vertically 4-connected matroids and cyclically 4-connected matroids, the analogues of 4-connected graphs and their duals. Nevertheless, chain theorems have been proved for certain classes of 3-connected matroids which are partially 4-connected. More precisely, instead of ruling out all 3-separations as one does in a 4-connected matroid, one can severely restrict the types of 3-separations that one allows. There are two natural ways of doing this. One way is to control the *structure* of 3-separations. A matroid is *sequentially 4-connected* if it is 3-connected and the only 3-separations (X, Y) are those such that, for some Z in $\{X, Y\}$, there is a *sequential ordering*, that is, an ordering (z_1, z_2, \dots, z_k) of Z such that $\{z_1, z_2, \dots, z_i\}$ is 3-separating for all i in $\{1, 2, \dots, k\}$. One raises connectivity to eliminate degeneracies and many of the degeneracies eliminated by requiring 4-connectivity are also eliminated by requiring sequential 4-connectivity. Geelen and Whittle [3] proved the following chain theorem.

Theorem 1.2. [3, Theorem 1.2] *Let M be a sequentially 4-connected matroid that is neither a wheel nor a whirl. Then M has an element z such that $M \setminus z$ or M/z is sequentially 4-connected.*

Another way to restrict 3-separations is to control *size*, that is, to require that they all have a small side. More precisely, let k be an integer exceeding one. A matroid M is $(4, k)$ -connected if M is 3-connected and, whenever (X, Y) is a 3-separating partition of $E(M)$, either $|X| \leq k$ or $|Y| \leq k$. Hall [4] called such a matroid *4-connected up to separators of size k* . Matroids that are $(4, 4)$ -connected have also been called *weakly 4-connected*. Rajan [9] showed that, for all positive integers m , a $(4, 4)$ -connected matroid M cannot be guaranteed to have a $(4, 4)$ -connected proper minor N with $|E(M)| - |E(N)| \leq m$. But Hall [4] proved that, by moving to $(4, 5)$ -connected matroids, one can get a chain theorem.

Theorem 1.3. *Let M be a $(4, 5)$ -connected matroid other than a rank-3 wheel. Then M has an element x such that $\text{co}(M \setminus x)$ or $\text{si}(M/x)$ is $(4, 5)$ -connected and has cardinality $|E(M)| - 1$ or $|E(M)| - 2$.*

In this paper, we prove a chain theorem where both the structure and the size of 3-separations is controlled, that is, where the allowable 3-separations are subject to both the restrictions imposed by Hall and those imposed by Geelen and Whittle. A 3-connected matroid M is $(4, k, S)$ -connected if M is both $(4, k)$ -connected and sequentially 4-connected.

Theorem 1.4. *Let M be a $(4, 5, S)$ -connected matroid that has no 5-element fans. Then M has an element x such that $M \setminus x$ or M/x is $(4, 5, S)$ -connected.*

Theorem 1.4 does not hold in certain highly structured matroids with 5-element fans. More generally, we have the following theorem, the main result of the paper.

Theorem 1.5. *Let M be a $(4, 5, S)$ -connected matroid other than a rank-3 wheel. Then M has an element x such that $\text{co}(M \setminus x)$ or $\text{si}(M/x)$ is $(4, 5, S)$ -connected and has cardinality $|E(M)| - 1$ or $|E(M)| - 2$.*

An example that illustrates the necessity of the 2-element move in Theorem 1.5 is given at the end of the paper. In proving this theorem, we shall use another new result, which seems to be of independent interest. A matroid that is $(4, 3)$ -connected is often called *internally 4-connected*.

Theorem 1.6. *Let M be a 4-connected matroid with $|E(M)| \geq 11$. Let $\{a, b, c, d, e\}$ be a rank-3 subset of $E(M)$. Then there are at least two elements x in $\{a, b, c, d, e\}$ such that $M \setminus x$ is internally 4-connected.*

The next section contains some basic definitions and results that will be needed in the proof of the main theorem. In Section 3, we outline how the proof of Theorem 1.5 proceeds. Basically, it divides the argument into the cases when M is $(4, k, S)$ -connected for $k = 2, 3, 4$, and 5 . Observe that M is $(4, 2, S)$ -connected if and only if it is 4-connected; and M is $(4, 3, S)$ -connected if and only if it is internally 4-connected. When M is 4-connected, there are two main cases to consider. The first uses Theorem 1.6, which is proved in Section 4; the second is treated in Section 5. The case when M is internally 4-connected is treated in Section 6. The proof of Theorem 1.5 is completed in Section 7 where the $(4, 4, S)$ -connected and $(4, 5, S)$ -connected cases are handled. The treatment of these cases is relatively short, but is somewhat artificially so since the latter relies crucially on Hall's proof of Theorem 1.3.

2. PRELIMINARIES

The matroid terminology used here will follow Oxley [5] except that the simplification and cosimplification of a matroid N will be denoted by $\text{si}(N)$ and $\text{co}(N)$, respectively. A *quad* in a matroid is a 4-element set that is both a circuit and a cocircuit. This paper will use some results and terminology from our papers describing the structure of 3-separations in 3-connected matroids [7, 8]. In this section, we introduce the relevant definitions. In addition, we prove some elementary connectivity results that will be used in the proof of the main theorem.

In a matroid M , a k -separating set X , or a k -separating partition $(X, E - X)$, or a k -separation $(X, E - X)$ is *exact* if $\lambda_M(X) = k - 1$. A k -separation $(X, E - X)$ is *minimal* if $|X| = k$ or $|E - X| = k$. It is well known (see, for example, [5, Corollary 8.1.11]) that if M is k -connected having $(X, E - X)$ as a k -separation with $|X| = k$, then X is a circuit or a cocircuit of M .

A set X in a matroid M is *fully closed* if it is closed in both M and M^* , that is, $\text{cl}(X) = X$ and $\text{cl}^*(X) = X$. The *full closure* of X , denoted $\text{fcl}(X)$, is the intersection of all fully closed sets that contain X . One way to obtain $\text{fcl}(X)$ is to take $\text{cl}(X)$, and then $\text{cl}^*(\text{cl}(X))$ and so on until neither the closure

nor coclosure operator adds any new elements of M . The full closure operator enables one to define a natural equivalence on exactly 3-separating partitions as follows. Two exactly 3-separating partitions (A_1, B_1) and (A_2, B_2) of a 3-connected matroid M are *equivalent*, written $(A_1, B_1) \cong (A_2, B_2)$, if $\text{fcl}(A_1) = \text{fcl}(A_2)$ and $\text{fcl}(B_1) = \text{fcl}(B_2)$. If $\text{fcl}(A_1) = E(M)$, then B_1 has a sequential ordering and we call B_1 *sequential*. Similarly, A_1 is sequential if $\text{fcl}(B_1) = E(M)$. We say (A_1, B_1) is *sequential* if A_1 or B_1 is sequential. Evidently, if $(A_1, B_1) \cong (A_2, B_2)$ and (A_1, B_1) is sequential, then so is (A_2, B_2) .

For a 3-connected matroid N , we shall be interested in 3-separations of N that show that it is not $(4, k, S)$ -connected. We call a 3-separation (X, Y) of N a $(4, k, S)$ -*violation* if either

- (i) $|X|, |Y| \geq k + 1$; or
- (ii) (X, Y) is non-sequential.

Observe that, when $k = 3$, condition (ii) implies condition (i). Hence (X, Y) is a $(4, 3, S)$ -violation of N if and only if $|X|, |Y| \geq 4$.

The following elementary lemma [7, Lemma 3.1] will be in repeated use throughout the paper.

Lemma 2.1. *For a positive integer k , let (A, B) be an exactly k -separating partition in a matroid M .*

- (i) *For e in $E(M)$, the partition $(A \cup e, B - e)$ is k -separating if and only if $e \in \text{cl}(A)$ or $e \in \text{cl}^*(A)$.*
- (ii) *For e in B , the partition $(A \cup e, B - e)$ is exactly k -separating if and only if e is in exactly one of $\text{cl}(A) \cap \text{cl}(B - e)$ and $\text{cl}^*(A) \cap \text{cl}^*(B - e)$.*
- (iii) *The elements of $\text{fcl}(A) - A$ can be ordered b_1, b_2, \dots, b_n so that $A \cup \{b_1, b_2, \dots, b_i\}$ is k -separating for all i in $\{1, 2, \dots, n\}$.*

The next well-known lemma specifies precisely when a single element z of a matroid M blocks a k -separating partition of $M \setminus z$ from extending to a k -separating partition of M . This result and its dual underlie numerous arguments in this paper.

Lemma 2.2. *In a matroid M with an element z , let (A, B) be a k -separating partition of $M \setminus z$. Then both $\lambda_M(A \cup z)$ and $\lambda_M(B \cup z)$ exceed $k - 1$ if and only if $z \in \text{cl}^*(A) \cap \text{cl}^*(B)$.*

Let S be a subset of a 3-connected matroid M . We call S a *fan* of M if $|S| \geq 3$ and there is an ordering (s_1, s_2, \dots, s_n) of the elements of S such that, for all i in $\{1, 2, \dots, n - 2\}$,

- (i) $\{s_i, s_{i+1}, s_{i+2}\}$ is a triangle or a triad; and
- (ii) when $\{s_i, s_{i+1}, s_{i+2}\}$ is a triangle, $\{s_{i+1}, s_{i+2}, s_{i+3}\}$ is a triad, and when $\{s_i, s_{i+1}, s_{i+2}\}$ is a triad, $\{s_{i+1}, s_{i+2}, s_{i+3}\}$ is a triangle.

The connectivity function λ_M of a matroid M has a number of attractive properties. For example, $\lambda_M(X) = \lambda_M(E - X)$. Moreover, the connectivity

functions of M and its dual M^* are equal. To see this, it suffices to note the easily verified fact that

$$\lambda_M(X) = r(X) + r^*(X) - |X|.$$

We shall often abbreviate λ_M as λ .

One of the most useful features of the connectivity function of M is that it is submodular, that is, for all $X, Y \subseteq E(M)$,

$$\lambda(X) + \lambda(Y) \geq \lambda(X \cap Y) + \lambda(X \cup Y).$$

This means that if X and Y are k -separating, and one of $X \cap Y$ or $X \cup Y$ is not $(k - 1)$ -separating, then the other must be k -separating. The next lemma specializes this fact.

Lemma 2.3. *Let M be a 3-connected matroid, and let X and Y be 3-separating subsets of $E(M)$.*

- (i) *If $|X \cap Y| \geq 2$, then $X \cup Y$ is 3-separating.*
- (ii) *If $|E(M) - (X \cup Y)| \geq 2$, then $X \cap Y$ is 3-separating.*

The last lemma will be in constant use throughout the paper. For convenience, we use the phrase *by uncrossing* to mean “by an application of Lemma 2.3.”

Another consequence of the submodularity of λ is the following very useful result for 3-connected matroids, which has come to be known as Bixby’s Lemma [1].

Lemma 2.4. *Let M be a 3-connected matroid and e be an element of M . Then either $M \setminus e$ or M/e has no non-minimal 2-separations. Moreover, in the first case, $\text{co}(M \setminus e)$ is 3-connected while, in the second case, $\text{si}(M/e)$ is 3-connected.*

A useful companion function to the connectivity function is the *local connectivity*, $\sqcap(X, Y)$, defined for sets X and Y in a matroid M , by

$$\sqcap(X, Y) = r(X) + r(Y) - r(X \cup Y).$$

Evidently,

$$\sqcap(X, E - X) = \lambda_M(X).$$

When M is \mathbb{F} -representable and hence viewable as a subset of the vector space $V(r(M), \mathbb{F})$, the local connectivity $\sqcap(X, Y)$ is precisely the rank of the intersection of those subspaces in $V(r(M), \mathbb{F})$ that are spanned by X and Y .

An attractive link between connectivity and local connectivity is provided by the next result [7, Lemma 2.6], which follows immediately by substitution.

Lemma 2.5. *Let X and Y be disjoint sets in a matroid M , then*

$$\lambda_M(X \cup Y) = \lambda_M(X) + \lambda_M(Y) - \sqcap_M(X, Y) - \sqcap_{M^*}(X, Y).$$

The first part of the next lemma [7, Lemma 2.3] is just a restatement of [5, Lemma 8.2.10]. The second part, which follows from the first, is the well-known fact that the connectivity function is monotone under taking minors.

Lemma 2.6. *Let M be a matroid.*

- (i) *Let X_1, X_2, Y_1 and Y_2 be subsets of $E(M)$. If $X_1 \subseteq Y_1$ and $X_2 \subseteq Y_2$, then $\cap(X_1, X_2) \leq \cap(Y_1, Y_2)$.*
- (ii) *If N is a minor of M and $X \subseteq E(M)$, then*

$$\lambda_N(X \cap E(N)) \leq \lambda_M(X).$$

One application of the last lemma that we shall use here is the following.

Lemma 2.7. *Let N be a 3-connected minor of a sequentially 4-connected matroid M . If (X, Y) is a 3-separation of M and $|X \cap E(N)|, |Y \cap E(N)| \geq 3$, then $(X \cap E(N), Y \cap E(N))$ is a sequential 3-separation of N .*

Proof. We may assume that X is sequential having (x_1, x_2, \dots, x_k) as a sequential ordering. Thus $(\{x_1, x_2, \dots, x_i\}, \{x_{i+1}, x_{i+2}, \dots, x_k\} \cup Y)$ is a 3-separation of M for all $i \geq 3$. We deduce that the lemma holds provided we can show that $(X \cap E(N), Y \cap E(N))$ is a 3-separation of N . But the latter follows immediately from Lemma 2.6. \square

The next lemma, which is elementary, is taken from Geelen and Whittle [3, Proposition 3.8].

Lemma 2.8. *Let M be a 3-connected matroid and (X, Y) be a non-sequential 3-separation of M . If $|X| = 4$, then X is a quad.*

In the next lemma, all but (ii) are taken from [3, Lemma 4.1]. The part of the lemma before (i) is in Coullard [2] (see also [5, Exercise 8.4.3]).

Lemma 2.9. *Let M be a 4-connected matroid and z be an element of M . Then $M \setminus z$ or M/z is weakly 4-connected. Let Q be a quad of M/z .*

- (i) *If (X, Y) is a 3-separation of $M \setminus z$ with $|X|, |Y| \geq 4$, then $|X \cap Q| = |Y \cap Q| = 2$.*
- (ii) *If T^* is a triad of $M \setminus z$ and $|E(M)| \geq 7$, then $Q \cap T^* \neq \emptyset$.*

Proof. (ii) Since M is 4-connected and $|E(M)| \geq 7$, the matroid M does not have Q as a quad or T^* as a triad. Thus $Q \cup z$ is a circuit of M and $T^* \cup z$ is a cocircuit of M . By orthogonality, $Q \cap T^* \neq \emptyset$. \square

The next lemma simplifies the task of identifying a $(4, 4, S)$ -violation.

Lemma 2.10. *Let N be a 3-connected matroid. Then (X, Y) is a $(4, 4, S)$ -violation if and only if*

- (i) $|X|, |Y| \geq 5$; or
- (ii) X and Y are non-sequential and at least one is a quad.

Proof. A 3-separation (X, Y) obeying (i) or (ii) is a $(4, 4, S)$ -violation. Conversely, suppose (X, Y) is a $(4, 4, S)$ -violation. We may assume that $|X|$ or $|Y|$ is at most 4. Then (X, Y) is non-sequential. Hence X and Y are non-sequential and at least one is a quad. \square

The notion of a flower was introduced in [7] to deal with crossing 3-separations, that is, 3-separations (A_1, A_2) and (B_1, B_2) for which each of the intersections $A_1 \cap B_1$, $A_1 \cap B_2$, $A_2 \cap B_1$, and $A_2 \cap B_2$ is non-empty. When each of these intersections has at least two elements, Lemma 2.3 implies that each is exactly 3-separating. Moreover, the union of any consecutive pair in the cyclic ordering $(A_1 \cap B_1, A_1 \cap B_2, A_2 \cap B_2, A_2 \cap B_1)$ is exactly 3-separating. This 4-tuple is an example of a flower.

An ordered partition (P_1, P_2, \dots, P_n) of the ground set of a 3-connected matroid M is a *flower* Φ if $\lambda_M(P_i) = 2 = \lambda_M(P_i \cap P_{i+1})$ for all i in $\{1, 2, \dots, n\}$ where all subscripts are interpreted modulo n . The sets P_1, P_2, \dots, P_n are the *petals* of Φ . It is shown in [7, Theorem 4.1] that every flower is either an *anemone* or a *daisy*. In the first case, all unions of petals are 3-separating; in the second, a union of petals is 3-separating if and only if the petals are consecutive in the cyclic ordering (P_1, P_2, \dots, P_n) . Observe that, when $n \leq 3$, the concepts of an anemone and a daisy coincide but, for $n \geq 4$, a flower cannot be both an anemone and a daisy.

Let Φ_1 and Φ_2 be flowers of a 3-connected matroid M . A natural quasi ordering on the collection of flowers of M is obtained by setting $\Phi_1 \preceq \Phi_2$ whenever every non-sequential 3-separation displayed by Φ_1 is equivalent to one displayed by Φ_2 . If $\Phi_1 \preceq \Phi_2$ and $\Phi_2 \preceq \Phi_1$, we say that Φ_1 and Φ_2 are *equivalent* flowers of M . Hence equivalent flowers display, up to equivalence of 3-separations, exactly the same non-sequential 3-separations of M . An element e of M is *loose* in a flower Φ if $e \in \text{fcl}(P_i) - P_i$ for some petal P_i of Φ .

The classes of anemones and daisies can be refined using local connectivity. For $n \geq 3$, an anemone (P_1, P_2, \dots, P_n) is called

- (i) a *paddle* if $\cap(P_i, P_j) = 2$ for all distinct i, j in $\{1, 2, \dots, n\}$;
- (ii) a *copaddle* if $\cap(P_i, P_j) = 0$ for all distinct i, j in $\{1, 2, \dots, n\}$; and
- (iii) *spike-like* if $n \geq 4$, and $\cap(P_i, P_j) = 1$ for all distinct i, j in $\{1, 2, \dots, n\}$.

Similarly, a daisy (P_1, P_2, \dots, P_n) is called

- (i) *swirl-like* if $n \geq 4$ and $\cap(P_i, P_j) = 1$ for all consecutive i and j , while $\cap(P_i, P_j) = 0$ for all non-consecutive i and j ; and
- (ii) *Vámos-like* if $n = 4$ and $\cap(P_i, P_j) = 1$ for all consecutive i and j , while $\{\cap(P_1, P_3), \cap(P_2, P_4)\} = \{0, 1\}$.

If (P_1, P_2, P_3) is a flower Φ and $\cap(P_i, P_j) = 1$ for all distinct i and j , we call Φ *ambiguous* if it has no loose elements, *spike-like* if there is an element in $\text{cl}(P_1) \cap \text{cl}(P_2) \cap \text{cl}(P_3)$ or $\text{cl}^*(P_1) \cap \text{cl}^*(P_2) \cap \text{cl}^*(P_3)$, and *swirl-like* otherwise. It is shown in [7] that every flower with at least three petals is one of these

six different *types*: a paddle, a copaddle, spike-like, swirl-like, Vámos-like, or ambiguous.

3. OUTLINE OF THE PROOF OF THE MAIN THEOREM

In this section, we begin by giving a slightly more detailed statement of the main theorem. Then we briefly outline the main steps in the proof of this theorem.

Theorem 3.1. *Let M be a $(4, 5, S)$ -connected matroid. Then M has an element x such that, for some N in $\{\text{co}(M \setminus x), \text{si}(M/x)\}$, the matroid N is $(4, 5, S)$ -connected. Moreover, $|E(N)| \in \{|E(M)| - 1, |E(M)| - 2, |E(M)| - 3\}$. In particular, $E(N) = |E(M)| - 3$ if and only if M is a rank-3 wheel; and $E(N) = |E(M)| - 1$ unless x is the element of a 5-element fan that is in two triangles or two triads of the fan.*

The overall strategy of the proof of this theorem is standard for proofs of theorems of this type. We begin by assuming that M is 4-connected. In that case, we prove the following result.

Theorem 3.2. *Let M be a 4-connected matroid with $|E(M)| \geq 13$. Then M has an element x such that $M \setminus x$ or M/x is $(4, 4, S)$ -connected.*

A crucial tool in this proof is the following result of Geelen and Whittle [3, Theorem 5.1].

Theorem 3.3. *Let M be a 4-connected matroid. Then M has an element z such that $M \setminus z$ or M/z is sequentially 4-connected.*

In proving Theorem 3.2, we have, by the last result and duality, that we may assume that the 4-connected matroid M has an element x for which $M \setminus x$ is sequentially 4-connected. If M is not $(4, 4, S)$ -connected, then it has a 3-separation (X, Y) with $|X|, |Y| \geq 5$. Moreover, this 3-separation is sequential. Hence it can be chosen so that $|X| = 5$ and X is sequential having $(x_1, x_2, x_3, x_4, x_5)$ as a sequential ordering. Because M is 4-connected, it has no triangles, so $\{x_1, x_2, x_3\}$ is a triad of $M \setminus x$. Now x_4 is in either the coclosure or the closure of $\{x_1, x_2, x_3\}$ in $M \setminus x$. In the first case, $\{x_1, x_2, x_3, x_4\}$ must be a union of triads in $M \setminus x$. Again, because M is 4-connected, it follows that every 4-element subset of $\{x_1, x_2, x_3, x_4, x\}$ is a cocircuit of M , that is, $M^*|_{\{x_1, x_2, x_3, x_4, x\}} \cong U_{3,5}$. The dual of this case is treated in Section 4 where Theorem 1.6 is proved. The second case, when $x_4 \in \text{cl}(\{x_1, x_2, x_3\})$, is treated in Section 5, thereby completing the proof of Theorem 3.2. That result imposed a lower bound on $|E(M)|$. By settling for a $(4, 5, S)$ -connected minor of M , we can drop this restriction. Specifically, at the end of Section 5, we prove the following result.

Corollary 3.4. *Let M be a 4-connected matroid. Then M has an element x such that $M \setminus x$ or M/x is $(4, 5, S)$ -connected.*

In view of the last result, when continuing the proof of Theorem 3.1 to the case when M is internally 4-connected, we may assume that M is not 4-connected. In that case, our proof uses the following result of Geelen and Whittle [3, Theorem 6.1].

Theorem 3.5. *Let T be a triangle in an internally 4-connected matroid M . Assume that M is not a wheel or whirl of rank three. Then either*

- (i) *T contains an element t for which $M \setminus t$ is sequentially 4-connected;*
or
- (ii) *$|E(M)| \leq 11$ and M has an element y such that M/y is sequentially 4-connected.*

Our main result in the internally 4-connected case is the following theorem, which is proved in Section 6.

Theorem 3.6. *Let M be a $(4, 3, S)$ -connected matroid that is not isomorphic to a wheel or whirl of rank three. Then M has an element e such that $M \setminus e$ or M/e is $(4, 5, S)$ -connected.*

The main difficulty in proving this theorem arises when $|E(M)|$ is relatively small although our argument does not differentiate cases based on $|E(M)|$.

The first theorem in Section 7 treats the case when M is $(4, 4, S)$ -connected by proving the following result.

Theorem 3.7. *Let M be a $(4, 4, S)$ -connected matroid that is not isomorphic to a wheel or whirl of rank 3 or 4. Then M has an element x such that $M \setminus x$ or M/x is $(4, 5, S)$ -connected.*

The core difficulties in proving this result have already been resolved in proving Theorem 3.6, so Theorem 3.7 has a short proof. By using the last result, we deduce that, to finish the proof of Theorem 3.1, we only need to treat the case when M is $(4, 5, S)$ -connected but not $(4, 4, S)$ -connected. This occupies the rest of Section 7. The proof here relies heavily on the detailed case analysis used by Hall in proving Theorem 1.3.

4. THE FIVE-POINT-PLANE CASE

In this section, we prove Theorem 1.6. It would be desirable to eliminate the lower bound on $|E(M)|$ in that theorem even though we do not need the stronger result to prove Theorem 1.5. To this end, the proof of Lemma 4.3 below includes more detail than is needed to get that result.

Lemma 4.1. *In a 4-connected matroid M , let $|F| = 5$ and $r(F) = 3$. For some $f \in F$, let (F_1, F_2) be a 3-separation of $M \setminus f$. Then*

- (i) *$|F_1 \cap F| = 2 = |F_2 \cap F|$; and*
- (ii) *if $|F_1| = 4$, then F_1 is a circuit of M and $F_1 \cup f$ contains a cocircuit of M containing f and having at least four elements.*

Proof. As M is 4-connected, exactly two elements of $F - f$ are in each of F_1 and F_2 , so (i) holds. Now let $|F_1| = 4$. Then $r_{M \setminus f}(F_1) + r_{M \setminus f}^*(F_1) - |F_1| = 2$, so $r_{M \setminus f}(F_1) + r_{M \setminus f}^*(F_1) = 6$. Since M has no triangles, $r(F_1) \geq 3$. Thus F_1 is a circuit unless $r(F_1) = 4$. In the exceptional case, $r_{M \setminus f}^*(F_1) = 2$, so every 3-element subset of F_1 is a triad in $M \setminus f$. Hence every 4-element subset of $F_1 \cup f$ is a cocircuit of M . Thus M has a 4-element cocircuit that contains exactly two elements of F . Since every 4-element subset of F is a circuit of M , we have a contradiction to orthogonality. We deduce that F_1 is indeed a circuit of M . Thus $r_{M \setminus f}^*(F_1) = 3$.

We now know that F_1 contains a cocircuit of $M \setminus f$. If this cocircuit is a triad T^* , then $T^* \cup f$ is a cocircuit of M containing f and contained in $F_1 \cup f$. We may now assume that F_1 is a cocircuit of $M \setminus f$. Since F_1 is not a quad of M , we deduce that $F_1 \cup f$ is a cocircuit of M . \square

Lemma 4.2. *In a 4-connected matroid M with $|E(M)| \geq 7$, let $\{a, b, c, d, e\}$ be a rank-3 subset of $E(M)$. Then*

- (i) $\text{co}(M \setminus a, b)$ is 3-connected;
- (ii) every non-trivial series class of $M \setminus a, b$ has exactly two elements and meets $\{c, d, e\}$; and
- (iii) each of c, d , and e is in at most one series pair of $M \setminus a, b$.

Proof. Consider $M \setminus a$. This matroid is certainly 3-connected. Now suppose that (X, Y) is a 2-separation of $M \setminus a, b$. Without loss of generality, we may assume that $\{d, e\} \subseteq X$. If $c \in X$, then $b \in \text{cl}(X)$ so $(X \cup b, Y)$ is a 2-separation of $M \setminus a$; a contradiction. Hence $c \in Y$. Again consider $(X \cup b, Y)$ and suppose that $|Y| \geq 3$. Then $(X \cup b, Y)$ is a 3-separation of $M \setminus a$ and $a \in \text{cl}(X \cup b)$, so $(X \cup b \cup a, Y)$ is a 3-separation of M ; a contradiction. Hence we may assume that $|Y| = 2$. Thus Y is a 2-cocircuit of $M \setminus a, b$ containing c . We deduce that $M \setminus a, b$ has no non-minimal 2-separations so $\text{co}(M \setminus a, b)$ is 3-connected. Moreover, every 2-cocircuit of $M \setminus a, b$ meets $\{c, d, e\}$. If both $\{c, y\}$ and $\{c, z\}$ are cocircuits of $M \setminus a, b$, then neither y nor z is in $\{d, e\}$, otherwise $\{a, b, c, d\}$ or $\{a, b, c, e\}$ is a quad of M ; a contradiction. Therefore $\{y, z\}$ is a cocircuit of $M \setminus a, b$ avoiding $\{c, d, e\}$. This contradiction implies that (ii) and (iii) hold. \square

Lemma 4.3. *In a 4-connected matroid M , let $r(\{a, b, c, d, e\}) = 3$. Suppose that (A_1, A_2) and (B_1, B_2) are 3-separations of $M \setminus a$ and $M \setminus b$, respectively, with $|A_1|, |A_2|, |B_1|, |B_2| \geq 4$ and $b \in A_1$ and $a \in B_1$. Then*

- (i) $\lambda_{M \setminus a, b}(A_1 \cap B_1) \in \{1, 2\}$;
- (ii) if $\lambda_{M \setminus a, b}(A_1 \cap B_1) = 1$ and $|E(M)| \geq 10$, then either $A_1 \cap B_1$ consists of a single element and this element is in $\{c, d, e\}$, or $A_1 \cap B_1$ consists of a 2-element cocircuit including exactly one element that is in $\{c, d, e\}$; in both cases, the two elements of $\{c, d, e\}$ that are not in $A_1 \cap B_1$ are in $A_2 \cap B_2$;

- (iii) if $\lambda_{M \setminus a, b}(A_1 \cap B_1) = 2$ and $|E(M)| \neq 10$, then $|A_2 \cap B_2| = 2$ and exactly one element of $\{c, d, e\}$ is in $A_1 \cap B_1$ while the other two elements are in $A_2 \cap B_2$, and $|A_2 \cap B_1| = |A_1 \cap B_2| = 2$.

Proof. Observe that, by orthogonality, we have:

4.3.1. Every cocircuit of M that meets $\{a, b, c, d, e\}$ does so in at least three elements.

Consider $M \setminus a, b$. From the preceding lemma, $\text{co}(M \setminus a, b)$ is 3-connected and each of c, d , and e is in at most one series pair of $M \setminus a, b$. Consider the placement of c, d , and e .

4.3.2. Either

- (I) exactly one element of $\{c, d, e\}$ is in each of $A_2 \cap B_1$, $A_2 \cap B_2$, and $A_1 \cap B_2$; or
- (II) exactly one element of $\{c, d, e\}$ is in $A_1 \cap B_1$ and the other two are in $A_2 \cap B_2$.

None of A_1, A_2, B_1 , and B_2 contains more than two elements of $\{a, b, c, d, e\}$. Since $a \in B_1$, exactly one of c, d , and e is in B_1 and the other two are in B_2 . Similarly, as $b \in A_1$, exactly one of c, d , and e is in A_1 and the other two are in A_2 .

Suppose that $|A_2 \cap B_1 \cap \{c, d, e\}| = 1$. Then, as $|B_1 \cap \{c, d, e\}| = 1$, we have $|A_1 \cap B_1 \cap \{c, d, e\}| = 0$. As $|A_1 \cap \{c, d, e\}| = 1$, it follows that $|A_1 \cap B_2 \cap \{c, d, e\}| = 1$. Since $|B_2 \cap \{c, d, e\}| = 2$, we deduce that $|A_2 \cap B_2 \cap \{c, d, e\}| = 1$. Hence if $|A_2 \cap B_1 \cap \{c, d, e\}| = 1$, then (I) holds. On the other hand, if $|A_2 \cap B_1 \cap \{c, d, e\}| = 0$, then $|A_1 \cap B_1 \cap \{c, d, e\}| = 1$, so $|A_2 \cap B_2 \cap \{c, d, e\}| = 2$ and (II) holds. This completes the proof of (4.3.2).

4.3.3. $\lambda_{M \setminus a, b}(A_2) = \lambda_{M \setminus a, b}(A_1 - b) = 2 = \lambda_{M \setminus a, b}(B_2) = \lambda_{M \setminus a, b}(B_1 - a)$.

By symmetry and taking complements, we see that it suffices to prove that $\lambda_{M \setminus a, b}(A_2) = 2$. Assume that $\lambda_{M \setminus a, b}(A_2) < 2$. Now $|A_1|, |A_2|, |B_1|, |B_2| \geq 4$, every series class of $M \setminus a, b$ has at most two elements and meets $\{c, d, e\}$ and $\text{co}(M \setminus a, b)$ is 3-connected. Thus, by (4.3.2), A_2 consists of exactly two series pairs each containing one member of $\{c, d, e\}$. Let these series pairs be $\{c, c'\}$ and $\{d, d'\}$. Since $|A_2| = 4$, by Lemma 4.1, A_2 is a circuit of M . But, in forming $\text{co}(M \setminus a, b)$, we contract one element from each of $\{c, c'\}$ and $\{d, d'\}$ to get a 2-element circuit. This contradicts the fact that $\text{co}(M \setminus a, b)$ is 3-connected since $|E(M)| \geq 9$. Hence (4.3.3) holds.

4.3.4. $b \in \text{cl}(A_1 - b)$ and $a \in \text{cl}(B_1 - a)$.

By symmetry, it suffices to prove that $b \in \text{cl}(A_1 - b)$. Assume the contrary. We have $r(A_1) + r(A_2) = r(M \setminus a) + 2$, so $r(A_1 - b) + r(A_2 \cup b) \leq r(M \setminus a) + 2$. Since $a \in \text{cl}(A_2 \cup b)$ and $|A_1 - b| \geq 3$, we deduce that $(A_1 - b, A_2 \cup b \cup a)$ is a 3-separation of M ; a contradiction. We conclude that (4.3.4) holds.

4.3.5. None of $A_1 \cap B_1$, $A_1 \cap B_2$, or $A_2 \cap B_1$ is empty.

If $A_1 \cap B_1 = \emptyset$, then $A_1 - b \subseteq B_2$, so, by (4.3.4), $b \in \text{cl}(B_2)$; a contradiction. If $A_1 \cap B_2 = \emptyset$, then $A_1 - b \subseteq B_1$, so $b \in \text{cl}(B_1)$; a contradiction. Hence $A_1 \cap B_2$ is non-empty and, by symmetry, so is $A_2 \cap B_1$.

4.3.6. *If $\lambda_{M \setminus a, b}(A_2 \cap B_2) \leq 2$, then $\lambda_{M \setminus a, b}(A_2 \cap B_2) = |A_2 \cap B_2|$.*

By (4.3.4), we deduce that $\lambda_{M \setminus a, b}(A_2 \cap B_2) = \lambda_{M \setminus a}(A_2 \cap B_2) = \lambda_M(A_2 \cap B_2)$. Since M is 4-connected, it follows that $\lambda_{M \setminus a, b}(A_2 \cap B_2) = |A_2 \cap B_2|$.

4.3.7. (i) $\lambda_{M \setminus a, b}(A_1 \cap B_2) = \lambda_{M \setminus b}(A_1 \cap B_2) = \lambda_M(A_1 \cap B_2)$; and
(ii) $\lambda_{M \setminus a, b}(A_2 \cap B_1) = \lambda_{M \setminus a}(A_2 \cap B_1) = \lambda_M(A_2 \cap B_1)$.

We have $|A_2 \cap \{c, d, e\}| = 2$ and $a \in \text{cl}(B_1 - a)$, so $\text{cl}((B_1 - a) \cup (A_2 \cap B_2))$ contains b . Thus (i) holds and (ii) follows by symmetry.

By submodularity, we have:

4.3.8. $\lambda_{M \setminus a, b}(A_1 \cap B_2) + \lambda_{M \setminus a, b}(A_2 \cap B_1) \leq 4$.

4.3.9. (i) *If $\lambda_{M \setminus a, b}(A_1 \cap B_2) \leq 2$, then $\lambda_{M \setminus a, b}(A_1 \cap B_2) = |A_1 \cap B_2|$.*
(ii) *If $\lambda_{M \setminus a, b}(A_2 \cap B_1) \leq 2$, then $\lambda_{M \setminus a, b}(A_2 \cap B_1) = |A_2 \cap B_1|$.*
(iii) *Either $|A_1 \cap B_2|$ or $|A_2 \cap B_1|$ is 1; or $|A_1 \cap B_2| = 2 = |A_2 \cap B_1|$.*
(iv) *If $|A_1 \cap B_2| = 1$, then A_1 is a 4-element circuit of M and $A_1 \cap B_1$ is a 2-element cocircuit of $M \setminus a, b$ that contains exactly one element of $\{c, d, e\}$.*

Parts (i) and (ii) follow from (4.3.7). Part (iii) follows by combining (i) and (ii) and using (4.3.8) and (4.3.5). To prove (iv), now assume that $|A_1 \cap B_2| = 1$. As $|A_2|, |A_1| \geq 4$, we have $|A_2 \cap B_2| \geq 3$ and $|A_1 \cap B_1| \geq 2$. Now $\lambda_{M \setminus a, b}(A_2 \cap B_2) = \lambda_{M \setminus a}(A_2 \cap B_2) \geq 3$, so $\lambda_{M \setminus a, b}(A_1 \cap B_1) \leq 1$. Hence $A_1 \cap B_1$ is a 2-element cocircuit of $M \setminus a, b$, so $|A_1| = 4$. Thus, by Lemma 4.1, A_1 is a circuit of M .

4.3.10. *Either*

- (i) $|A_2 \cap B_2| = 2$ and $\lambda_{M \setminus a, b}(A_1 \cap B_1) \leq 2$; or
- (ii) $|A_2 \cap B_2| \geq 3$ and $\lambda_{M \setminus a, b}(A_1 \cap B_1) = 1$.

Moreover, if $\lambda_{M \setminus a, b}(A_1 \cap B_1) = 1$, then either $|A_1 \cap B_1| = 1$, or $A_1 \cap B_1$ is a 2-cocircuit of $M \setminus a, b$ that contains exactly one element of $\{c, d, e\}$.

To see this, note that, by (4.3.9)(iii), either $|A_1 \cap B_2| = |A_2 \cap B_1| = 2$; or $|A_1 \cap B_2|$ or $|A_2 \cap B_1|$ is 1. Thus, as $|B_2|, |A_2| \geq 4$, we have $|A_2 \cap B_2| \geq 2$. Also $\lambda_{M \setminus a, b}(A_2 \cap B_2) = \lambda_{M \setminus a}(A_2 \cap B_2)$ as $b \in \text{cl}(A_1 - b)$. Hence $\lambda_{M \setminus a, b}(A_2 \cap B_2) \geq 2$, so, by submodularity, $\lambda_{M \setminus a, b}(A_1 \cap B_1) \leq 2$. Moreover, if $|A_2 \cap B_2| \geq 3$, then $\lambda_{M \setminus a, b}(A_2 \cap B_2) \geq 3$, so $\lambda_{M \setminus a, b}(A_1 \cap B_1) \leq 1$. We deduce that (i) or (ii) of (4.3.10) holds. The final assertion of the sublemma follows directly from Lemma 4.2.

By (4.3.5) and (4.3.10), we deduce that (i) of the lemma holds.

4.3.11. *If $|A_1 \cap B_1| = 1$ and $|E(M)| \geq 10$, then $A_1 \cap B_1 \subseteq \{c, d, e\}$.*

By (ii) of the lemma, each of A_1 and B_1 has exactly four elements. By Lemma 4.1, each of A_1 and B_1 is a circuit and $A_1 \cup a$ and $B_1 \cup b$ contain cocircuits C_a^* and C_b^* of M containing a and b , respectively. As $|A_1 \cap B_1| = 1$ and each of these cocircuits contains at least four elements, C_a^* and C_b^* are distinct.

Assume that (4.3.11) fails. Then (I) of (4.3.2) holds and $|A_1 \cap B_2 \cap \{c, d, e\}| = 1 = |A_2 \cap B_1 \cap \{c, d, e\}|$. Let $A_1 \cap B_2 = \{c, x\}$, let $A_2 \cap B_1 = \{d, y\}$, and let $A_1 \cap B_1 = \{z\}$. Then $A_1 \cup B_1$ is spanned by $\{a, b, c, z\}$ since we have the circuits $\{b, c, z, x\}$, $\{a, b, c, d\}$, and $\{a, d, z, y\}$. Thus

$$\begin{aligned} \lambda(A_1 \cup B_1) &= r(A_1 \cup B_1) + r^*(A_1 \cup B_1) - |A_1 \cup B_1| \\ &\leq 4 + 5 - 7 = 2. \end{aligned}$$

This contradicts the fact that M is 4-connected because $|A_2 \cap B_2| \geq 3$ since $|E(M)| \geq 10$. We conclude that (4.3.11) holds.

By combining (4.3.10) and (4.3.11), we deduce that (ii) of the lemma holds.

As an immediate consequence of (4.3.10), we have:

4.3.12. *If $\lambda_{M \setminus a, b}(A_1 \cap B_1) = 2$, then $|A_2 \cap B_2| = 2$.*

We now complete the proof of (iii) of the lemma. Assume that $\lambda_{M \setminus a, b}(A_1 \cap B_1) = 2$. Then, by (4.3.12), $|A_2 \cap B_2| = 2$. Since $|A_2|, |B_2| \geq 4$, it follows by (4.3.9)(iii) that $|A_1 \cap B_2| = |A_2 \cap B_1| = 2$. Suppose that (I) of (4.3.2) holds. Then $\{a, b\} \subseteq \text{cl}(E - \{a, b\} - (A_1 \cap B_1))$, so $\lambda_{M \setminus a, b}(A_1 \cap B_1) = 2 = \lambda_{M \setminus a}(A_1 \cap B_1) = \lambda_M(A_1 \cap B_1)$. Hence, as $|A_2| \geq 4$, we deduce that $|A_1 \cap B_1| = 2$ and, therefore, $|E(M)| = 10$. Thus, provided $|E(M)| \neq 10$, we may assume that (II) of (4.3.2) holds and part (iii) of the lemma follows. \square

The essential fact from the last lemma needed for the Proof of Theorem 1.6 is the following.

Corollary 4.4. *In a 4-connected matroid M with $|E(M)| \geq 11$, let $r(\{a, b, c, d, e\}) = 3$. Suppose that (A_1, A_2) and (B_1, B_2) are 3-separations of $M \setminus a$ and $M \setminus b$, respectively, with $|A_1|, |A_2|, |B_1|, |B_2| \geq 4$ and $b \in A_1$ and $a \in B_1$. Then one element of $\{c, d, e\}$ is in $A_1 \cap B_1$ and the other two are in $A_2 \cap B_2$.*

Proof of Theorem 1.6. Suppose that none of $M \setminus a, M \setminus b, M \setminus c$, and $M \setminus d$ is internally 4-connected. Let $(A_1, A_2), (B_1, B_2), (C_1, C_2)$, and (D_1, D_2) be 3-separations of $M \setminus a, M \setminus b, M \setminus c$, and $M \setminus d$ with $|A_1|, |A_2|, |B_1|, |B_2|, |C_1|, |C_2|, |D_1|, |D_2| \geq 4$. Each of the last eight sets contains exactly two elements of $\{a, b, c, d, e\}$. In particular, we may assume that $\{b, c\} \subseteq A_1 \cap \{b, c, d, e\}$. Label B_1 and C_1 so that $a \in B_1 \cap C_1$. By Corollary 4.4, since $|A_1 \cap B_1 \cap \{c, d, e\}| = 1$, we deduce that $c \in B_1$, so $B_2 \cap \{a, c, d, e\} = \{d, e\}$. Symmetrically, $b \in C_1$.

Now consider (D_1, D_2) labelling this so that $a \in D_1$. Because $d \in A_2$ and $A_2 \cap \{b, c, d, e\} = \{d, e\}$, we deduce that $D_1 \cap A_2 \cap \{a, b, c, d, e\} = \{e\}$.

Thus $D_1 \cap \{a, b, c, d, e\} = \{a, e\}$ and $D_2 \cap \{a, b, c, d, e\} = \{b, c\}$. Now $d \in B_2$ and $b \in D_2$, yet $D_2 \cap B_2 \cap \{a, b, c, d, e\} = \emptyset$. This contradiction to Corollary 4.4 completes the proof that at least one of $M \setminus a$, $M \setminus b$, $M \setminus c$, and $M \setminus d$ is internally 4-connected. If exactly one of $M \setminus a$, $M \setminus b$, $M \setminus c$, and $M \setminus d$ is internally 4-connected, assume it is $M \setminus a$. Then, arguing as above, we get that at least one of $M \setminus b$, $M \setminus c$, $M \setminus d$, and $M \setminus e$ is internally 4-connected. We conclude that at least two of $M \setminus a$, $M \setminus b$, $M \setminus c$, $M \setminus d$, and $M \setminus e$ are internally 4-connected. \square

5. THE 4-CONNECTED CASE

In this section, we shall complete the proof of Theorem 3.2, thereby proving the main theorem in the case that M is 4-connected. We are following the strategy outlined in Section 3. The key remaining result we need to get Theorem 3.2 is the following.

Theorem 5.1. *Let M be a 4-connected matroid with $|E(M)| \geq 13$. Let x be an element of M such that $M \setminus x$ is sequentially 4-connected but not weakly 4-connected, and M/x is not sequentially 4-connected. Suppose that $\{s, t, u\}$ is a triad of $M \setminus x$, that $\{s, t, u, y\}$ is a circuit of $M \setminus x$, and that $\{s, t, u, y, c\}$ is 3-separating in $M \setminus x$. Then, for some z in $\{s, t, u\}$, the matroid M/z is $(4, 4, S)$ -connected.*

Proof. Since $M \setminus x$ is not weakly 4-connected, by Lemma 2.9, we have:

5.1.1. M/x is weakly 4-connected.

Since M/x is not sequentially 4-connected, by Lemma 2.8,

5.1.2. M/x has a quad D .

Assume the theorem fails.

Lemma 5.2. *The matroid M/s has a $(4, 4, S)$ -violator (S_1, S_2) with $\{t, u, y\} \subseteq S_1$ and x in S_2 .*

Proof. Because the theorem fails, M/s has a $(4, 4, S)$ -violator (S_1, S_2) where we can label this so that $|S_1 \cap \{t, u, y\}| \geq 2$.

5.2.1. *If $\{t, u, y\} \subseteq S_1$, then $x \in S_2$.*

To see this, assume that $x \in S_1$. We have

$$r_{M/s}(S_1) + r_{M/s}(S_2) = r(M/s) + 2,$$

so $r(S_1 \cup s) + r(S_2 \cup s) = r(M) + 3$. But $\{s, t, u, x\}$ is a cocircuit of M and $\{t, u, x\} \cap S_2 = \emptyset$. Hence $r(S_2 \cup s) = r(S_2) + 1$. Thus $(S_1 \cup s, S_2)$ is a 3-separation of M ; a contradiction. Hence (5.2.1) holds.

We may now assume that $|S_1 \cap \{t, u, y\}| = 2$. Then $(S_1 \cup \{t, u, y\}, S_2 - \{t, u, y\})$ is a 3-separation of M/s that is equivalent to (S_1, S_2) . Hence S_2 is not a quad of M/s . Thus $(S_1 \cup \{t, u, y\}, S_2 - \{t, u, y\})$ is a $(4, 4, S)$ -violator unless $|S_2| = 5$ and $S_2 - \{t, u, y\}$ is not a quad of M/s . We deduce

that the lemma holds unless S_2 is a sequential 3-separating set of M/s having a sequential ordering $(1, 2, 3, 4, 5)$ with $5 \in \{t, u, y\}$.

Consider the exceptional case. As $5 \in \{t, u, y\}$, we have $5 \in \text{cl}_{M/s}(S_1)$. Thus $5 \in \text{cl}_{M/s}(\{1, 2, 3, 4\})$. Since M/s has no triads, we deduce that $\{1, 2, 3\}$ is a triangle of M/s . If $4 \in \text{cl}_{M/s}(\{1, 2, 3\})$, then $M|_{\{1, 2, 3, 4, s\}} \cong U_{3,5}$. By applying the argument for (5.2.1) to $(S_1 \cup \{t, u, y\}, S_2 - \{t, u, y\})$, we deduce that $x \in \{1, 2, 3, 4\}$. But this means that the circuit $\{1, 2, 3, 4\}$ meets the cocircuit $\{s, t, u, x\}$ in a single element; a contradiction. Hence $4 \notin \text{cl}_{M/s}(\{1, 2, 3\})$, so $\{1, 2, 3, 4\}$ is a cocircuit of M/s and hence of M . Moreover, $\{1, 2, 3, s\}$ is a circuit of M . By orthogonality, $x \in \{1, 2, 3\}$ so, since 1, 2, and 3 can be arbitrarily reordered, we may assume that $x = 1$.

Let $Z = \{x, 2, 3, 4, s, t, u, y\}$. Then $r_{M/s}(Z - s) \leq 4$ since $Z - s$ is spanned in M/s by $\{2, 3, 4\}$ together with an element of $\{t, u, y\} - 5$ because $5 \in \text{cl}_{M/s}(\{1, 2, 3, 4\})$ and $5 \in \{t, u, y\}$. Now $\{s, t, u, y\}$ is 3-separating in $M \setminus x$. Thus, by Lemma 2.9, the quad D of M/x satisfies

$$|D \cap \{s, t, u, y\}| = 2 \text{ and } |D - \{s, t, u, y\}| = 2.$$

Now D is a cocircuit of M and $D \cup x$ is a circuit of M . As the cocircuit $\{x, 2, 3, 4\}$ meets $D \cup x$, orthogonality implies that D meets $\{2, 3, 4\}$.

We now have two possibilities:

- (i) $D \subseteq Z$; and
- (ii) $D - Z = \{d\}$ for some element d .

In the first case, D contains exactly two elements of $\{2, 3, 4\}$. Consider M^* . It has $\{x, 2, 3, 4\}$, $\{s, t, u, x\}$, and D among its circuits. Let B^* consist of $\{x, y\}$ together with two elements of $\{s, t, u\}$, both in D if possible, and one element of $\{2, 3, 4\} \cap D$. Then B^* spans Z in M^* . Hence $r_{M^*}(Z) \leq 5$. But we have already shown that $r_M(Z) \leq 5$. Thus $r_M(Z) + r_{M^*}(Z) - |Z| \leq 2$, so $|E(M) - Z| \leq 2$. Hence $|E(M)| \leq 10$; a contradiction.

In case (ii), the circuit $D \cup x$ and the fact that $r(Z) \leq 5$ imply that $r(Z \cup d) \leq 5$. Moreover, $Z \cup d$ is spanned in M^* by $\{x, y, s, t, 2, 3\}$, so $r^*(Z \cup d) \leq 6$. Thus $r_M(Z \cup d) + r_{M^*}(Z \cup d) - |Z \cup d| \leq 2$, so $|E(M) - (Z \cup d)| \leq 2$. Hence $|E(M)| \leq 11$. This contradiction completes the proof of the lemma. \square

Lemma 5.3. *If (S_1, S_2) is a $(4, 4, S)$ -violator of M/s with $\{t, u, y\} \subseteq S_1$ and $x \in S_2$, then*

- (i) $r_{M/s}(S_1), r_{M/s}(S_2) \geq 3$; and
- (ii) *either $|S_1|, |S_2| \geq 5$, or S_2 is a quad of M/s and S_1 is non-sequential but is not a quad.*

Proof. Suppose that $r_{M/s}(S_2) = 2$. Then, by Lemma 2.10, $|S_2| \geq 5$ and so every 4-element subset of S_2 is a circuit of M . Thus M has a 4-element circuit meeting the cocircuit $\{s, t, u, x\}$ in $\{x\}$. This contradicts orthogonality. Thus $r_{M/s}(S_2) \geq 3$.

Now assume that $r_{M/s}(S_1) = 2$. Then, by Lemma 2.10, $|S_1| \geq 5$. Now take a and b to be distinct elements of $S_1 - \{t, u, y\}$. Then $\{a, b, y, s\}$ is a

circuit of M meeting the cocircuit $\{s, t, u, x\}$ in a single element; a contradiction to orthogonality. We conclude that (i) holds.

To prove (ii), note that if it fails, then S_1 is a quad of M/s . But S_1 is not a quad of the 4-connected matroid M , so $S_1 \cup s$ is a circuit of M that properly contains the circuit $\{s, t, u, y\}$; a contradiction. \square

Now, by Lemma 5.2, we can choose (S_1, S_2) , (T_1, T_2) , and (U_1, U_2) to be $(4, 4, S)$ -violators of M/s , M/t , and M/u , respectively, with $x \in S_2 \cap T_2 \cap U_2$ and $(S_2 \cup T_2 \cup U_2) \cap \{s, t, u, y\} = \emptyset$. Let S'_2, T'_2 , and U'_2 be $S_2 - x, T_2 - x$, and $U_2 - x$, respectively. In the results that follow, we prove various properties of the set S_2 . By symmetry, the corresponding properties will also hold for T_2 and U_2 .

Lemma 5.4. *The elements s and x are in $\text{cl}_M(S_2)$ and $\text{cl}_{M/s}(S'_2)$, respectively. Thus $x \in \text{cl}_M(S'_2 \cup s)$.*

Proof. We have

$$r_{M/s}(S_1) + r_{M/s}(S_2) = r(M/s) + 2.$$

Assume $x \notin \text{cl}_{M/s}(S'_2)$. Then

$$r_{M/s}(S_1 \cup x) + r_{M/s}(S'_2) = r(M/s) + 2,$$

so $r(S_1 \cup x \cup s) + r(S'_2 \cup s) = r(M) + 3$. Now $\{s, t, u, x\}$ is a cocircuit of M meeting $S'_2 \cup s$ in a single element. Hence $r(S'_2 \cup s) = r(S'_2) + 1$. Thus $r(S_1 \cup x \cup s) + r(S'_2) = r(M) + 2$. But M is 4-connected, so $|S'_2| \leq 2$. This contradicts the fact that $|S_2| \geq 4$. We deduce that $x \in \text{cl}_{M/s}(S'_2)$. Hence $x \in \text{cl}_M(S'_2 \cup s)$. But $x \notin \text{cl}_M(S'_2)$ because $\{s, t, u, x\}$ is a cocircuit that avoids S'_2 . Hence $s \in \text{cl}_M(S'_2 \cup x) = \text{cl}(S_2)$. \square

Lemma 5.5. $\Pi(\{s, t, u, y\}, S'_2 \cup T'_2 \cup U'_2) = 2$.

Proof. The set $\{s, t, u, y\}$ is 3-separating in $M \setminus x$, so $\Pi(\{s, t, u, y\}, E(M) - \{s, t, u, y, x\}) = 2$. By Lemma 2.6(i), $\Pi(\{s, t, u, y\}, S'_2 \cup T'_2 \cup U'_2) \leq 2$.

Now $r(\{s, t, u, y\}) = 3$ and, by Lemma 5.4 and symmetry, $\text{cl}(S'_2 \cup T'_2 \cup U'_2 \cup x)$ contains $\{s, t, u\}$ and hence y . Thus

$$3 = r(\{s, t, u, y\}) + r(S'_2 \cup T'_2 \cup U'_2 \cup x) - r(S'_2 \cup T'_2 \cup U'_2 \cup x \cup \{s, t, u, y\})$$

and

$$\begin{aligned} 2 &\geq r(\{s, t, u, y\}) + r(S'_2 \cup T'_2 \cup U'_2) - r(S'_2 \cup T'_2 \cup U'_2 \cup \{s, t, u, y\}) \\ &\geq r(\{s, t, u, y\}) + r(S'_2 \cup T'_2 \cup U'_2) - r(S'_2 \cup T'_2 \cup U'_2 \cup x \cup \{s, t, u, y\}) \\ &\geq r(\{s, t, u, y\}) + r(S'_2 \cup T'_2 \cup U'_2 \cup x) - 1 \\ &\quad - r(S'_2 \cup T'_2 \cup U'_2 \cup x \cup \{s, t, u, y\}) \\ &= 3 - 1 = 2. \end{aligned}$$

We conclude that $\Pi(\{s, t, u, y\}, S'_2 \cup T'_2 \cup U'_2) = 2$. \square

Lemma 5.6. *If $\lambda_{M \setminus x}(S'_2 \cup T'_2 \cup U'_2) = 2$, then*

$$E(M) = S'_2 \cup T'_2 \cup U'_2 \cup \{s, t, u, y, x\}.$$

Proof. By Lemma 2.5, we have

$$\begin{aligned} \lambda_{M \setminus x}(S'_2 \cup T'_2 \cup U'_2 \cup \{s, t, u, y\}) &\leq \lambda_{M \setminus x}(S'_2 \cup T'_2 \cup U'_2) + \lambda_{M \setminus x}(\{s, t, u, y\}) \\ &\quad - \cap(S'_2 \cup T'_2 \cup U'_2, \{s, t, u, y\}) \\ &= 2 + 2 - 2 = 2. \end{aligned}$$

But $x \in \text{cl}(S'_2 \cup T'_2 \cup U'_2 \cup \{s, t, u, y\})$, so $\lambda_M(S'_2 \cup T'_2 \cup U'_2 \cup \{s, t, u, y, x\}) \leq 2$.

The matroid M is 4-connected, so $E(M) - (S'_2 \cup T'_2 \cup U'_2 \cup \{s, t, u, y, x\})$ is a set V with at most two elements. To complete the proof of the lemma, we need to show that V is empty.

First we show that

5.6.1. $V \subseteq \text{cl}(\{s, t, u\})$.

Assume not. As $\lambda_{M \setminus x}(S'_2 \cup T'_2 \cup U'_2) = 2$, we have

$$\begin{aligned} 2 &= r(S'_2 \cup T'_2 \cup U'_2) + r(\{s, t, u, y\} \cup V) - r(M \setminus x) \\ &\geq r(S'_2 \cup T'_2 \cup U'_2) + r(\{s, t, u, y\}) - r(M \setminus x) \\ &= 2 \end{aligned}$$

where the last step holds by Lemma 5.5 since $r(M \setminus x) = r(M \setminus x \setminus V)$ as $|V \cup x| \leq 3$. Thus equality holds throughout the last chain of inequalities, so $V \subseteq \text{cl}(\{s, t, u, y\}) = \text{cl}(\{s, t, u\})$, that is, (5.6.1) holds.

Now take $e \in V$. Then $\{s, t, u, e\}$ and $\{s, t, u, y\}$ are both circuits of M , so every 4-element subset of $\{s, t, u, y, e\}$ is a circuit of M . By (5.1.2), M/x has a quad D . By Lemma 2.9, D contains exactly two elements of $\{s, t, u, y, e\}$. But this contradicts orthogonality since D is a cocircuit of M . We conclude that $V = \emptyset$. Hence the lemma holds. \square

Lemma 5.7. *The matroid $M \setminus x/s$ is 3-connected.*

Proof. Certainly $M \setminus x$ is 3-connected and has no triangles since M is 4-connected. The matroid $M \setminus x/s$ has $\{t, u, y\}$ as a triangle and is simple and cosimple. Assume (X, Y) is a 2-separation of $M \setminus x/s$. Since $M \setminus x/s$ has no 2-cocircuits, this 2-separation is non-minimal. Then, without loss of generality, $|X \cap \{t, u, y\}| \geq 2$. Therefore $(X \cup \{t, u, y\}, Y - \{t, u, y\})$ is a 2-separation of $M \setminus x/s$ and $|Y - \{t, u, y\}| \geq 3$. Hence we may assume that $X \supseteq \{t, u, y\}$ and $|Y| \geq 3$. Now

$$r_{M \setminus x/s}(X) + r_{M \setminus x/s}(Y) = r(M \setminus x/s) + 1.$$

So $r(X \cup s) + r(Y \cup s) = r(M) + 2$. We have $\{s, t, u, x\}$ as a cocircuit of M , so $\{s, t, u\}$ is a cocircuit of $M \setminus x$. Hence, as $\{t, u\} \subseteq X$, we have $r(Y \cup s) = r(Y) + 1$, so $r(X \cup s) + r(Y) = r(M) + 1$. Thus $r(X \cup s) + r(Y \cup x) \leq r(M) + 2$, a contradiction to the fact that M is 4-connected. \square

Lemma 5.8. *The partition $(S_1 \cup s, S'_2)$ is a vertical 3-separation of $M \setminus x$, so $\lambda_{M \setminus x}(S'_2) = 2$. Moreover, if $|S'_2| = 3$, then S'_2 is a triad of $M \setminus x$.*

Proof. We have

$$\begin{aligned}
 r(S_1 \cup s) + r(S'_2) - r(M \setminus x) &= [r_{M/s}(S_1) + 1] + [r(S'_2 \cup s) - 1] \\
 &\quad - [r(M/s) + 1] \\
 &= r_{M/s}(S_1) + r_{M/s}(S'_2) - r(M/s) \\
 &= r_{M/s}(S_1) + r_{M/s}(S_2) - r(M/s) \\
 &= 2.
 \end{aligned}$$

Thus $(S_1 \cup s, S'_2)$ is a 3-separation of $M \setminus x$. Since, by Lemmas 5.3 and 5.4, $r_{M/s}(S_1) \geq 3$ and $r_{M/s}(S'_2) = r_{M/s}(S_2) \geq 3$, it follows that this 3-separation is vertical.

Finally, if $|S'_2| = 3$, then $(S_1 \cup x, S'_2)$ is a minimal 3-separation of $M \setminus x$. As $M \setminus x$ has no triangles, it follows that S'_2 is a triad of $M \setminus x$. \square

Lemma 5.9. $S'_2 \cap T'_2 \neq \emptyset$.

Proof. Assume $S'_2 \cap T'_2 = \emptyset$. Then $S'_2 \subseteq T_1$ and $s \in T_1$. But, by Lemma 5.4, $x \in \text{cl}(S'_2 \cup s)$. Hence $x \in \text{cl}(T_1)$. Thus, by Lemma 5.8 and symmetry, $(T_1 \cup t \cup x, T'_2)$ is a 3-separation of M ; a contradiction. \square

Lemma 5.10. *The sets S'_2 and T'_2 have the following properties:*

- (i) $\lambda_M(S'_2 - T'_2) + \lambda_M(T'_2 - S'_2) \leq 4$;
- (ii) if $|S'_2 - T'_2| \geq 2$, then $|T'_2 - S'_2| \leq 2$;
- (iii) if $|S'_2 - T'_2| \geq 3$, then $|T'_2 - S'_2| \leq 1$; and
- (iv) if $|S'_2 - T'_2|, |T'_2 - S'_2| \geq 2$, then $|S'_2 - T'_2| = |T'_2 - S'_2| = 2$.

Proof. We have $\lambda_{M \setminus x}(S'_2) = 2 = \lambda_{M \setminus x}(T'_2)$ while $E(M \setminus x) - S'_2 = S_1 \cup s$ and $E(M \setminus x) - T'_2 = T_1 \cup t$. Thus

$$\begin{aligned}
 4 &= \lambda_{M \setminus x}(S'_2) + \lambda_{M \setminus x}(T_1 \cup t) \\
 &\geq \lambda_{M \setminus x}(S'_2 \cup T_1 \cup t) + \lambda_{M \setminus x}(S'_2 \cap (T_1 \cup t)) \\
 &= \lambda_{M \setminus x}(T'_2 - S'_2) + \lambda_{M \setminus x}(S'_2 - T'_2) \\
 &= \lambda_M(T'_2 - S'_2) + \lambda_M(S'_2 - T'_2).
 \end{aligned}$$

The last step here holds because $E(M \setminus x) - (T'_2 - S'_2) \supseteq S'_2 \cup s$ and $x \in \text{cl}_M(S'_2 \cup s)$, so $\lambda_{M \setminus x}(T'_2 - S'_2) = \lambda_M(T'_2 - S'_2)$ and, by symmetry, $\lambda_{M \setminus x}(S'_2 - T'_2) = \lambda_M(S'_2 - T'_2)$. Thus (i) holds. Since M is 4-connected, parts (ii) and (iii) hold. Part (iv) follows by using (ii) and the natural symmetric form of it. \square

Lemma 5.11. *If $|S'_2 \cap T'_2| = 1$, then S'_2 and T'_2 are both triads of $M \setminus x$.*

Proof. Suppose that S'_2 is not a triad of $M \setminus x$. Then, by Lemma 5.8, $|S'_2| > 3$, so $|S'_2 - T'_2| \geq 3$. Hence, by Lemma 5.10(iii), $|T'_2 - S'_2| \leq 1$. As $|T'_2 \cap S'_2| = 1$, it follows that $|T'_2| \leq 2$; a contradiction. We conclude that S'_2 is a triad and, by symmetry, so is T'_2 . \square

Lemma 5.12. *If each of $S'_2 - T'_2$, $T'_2 - S'_2$, and $S'_2 \cap T'_2$ has at least two elements, then $(S'_2 \cap T'_2, T'_2 - S'_2, (S_1 \cup s) \cap (T_1 \cup t), S'_2 - T'_2)$ is a Vámos-like flower Φ in $M \setminus x$ and $|S'_2 - T'_2| = 2 = |T'_2 - S'_2|$.*

Proof. By Lemma 5.10, we deduce that each of $S'_2 - T'_2$ and $T'_2 - S'_2$ has exactly two elements and so is 3-separating in $M \setminus x$. We have $\lambda_{M \setminus x}(S'_2) = 2 = \lambda_{M \setminus x}(T'_2)$ while $|(S_1 \cup s) \cap (T_1 \cup t)| = |E(M \setminus x) - (S'_2 \cup T'_2)| \geq |\{s, t, u, y\}| \geq 4$. We deduce, by Lemma 2.3, that $\lambda_{M \setminus x}(S'_2 \cap T'_2) = 2 = \lambda_{M \setminus x}((S_1 \cup s) \cap (T_1 \cup t))$. Hence Φ is a flower in $M \setminus x$. Now $(S_1 \cup s) \cap (T_1 \cup t)$ is 3-separating in $M \setminus x$ and has at least four elements. Thus, by Lemma 2.9, D has exactly two elements in $(S_1 \cup s) \cap (T_1 \cup t)$. Similarly, D has exactly two elements in S'_2 and exactly two elements in T'_2 . Hence D has exactly two elements in $S'_2 \cap T'_2$. We deduce, since D contains a cocircuit of $M \setminus x$, that $\cap_{M \setminus x}^*(S'_2 \cup T'_2, (S_1 \cup s) \cap (T_1 \cup t)) > 0$.

Now D avoids the 4-element set $(S'_2 - T'_2) \cup (T'_2 - S'_2)$ of $E(M \setminus x)$ so, by Lemma 2.9 again, the set $(S'_2 - T'_2) \cup (T'_2 - S'_2)$ is not exactly 3-separating. Thus Φ is a daisy in each of $M \setminus x$ and $(M \setminus x)^*$. As $\cap_{M \setminus x}^*(S'_2 \cup T'_2, (S_1 \cup s) \cap (T_1 \cup t)) > 0$, the flower Φ is not swirl-like in $(M \setminus x)^*$. Hence Φ is not swirl-like in $M \setminus x$, so Φ is Vámos-like. \square

Lemma 5.13. *If $|S'_2 \cap T'_2| \geq 2$, then $|S'_2 - T'_2| \leq 1$ or $|T'_2 - S'_2| \leq 1$.*

Proof. Assume that both $S'_2 - T'_2$ and $T'_2 - S'_2$ exceed one. Then, by Lemma 5.12, Φ is a Vámos-like flower in $M \setminus x$ and $|S'_2 - T'_2| = 2 = |T'_2 - S'_2|$. By [7, Theorem 6.1], Φ has no loose elements.

Now $(S'_2 - T'_2) \cup [(S_1 \cup s) \cap (T_1 \cup t)] = T_1 \cup t$ and $(T'_2, T_1 \cup t)$ is a 3-separation of $M \setminus x$. Hence it is sequential. Assume that $T_1 \cup t$ is sequential and consider the set F of the first three elements in a sequential ordering $\overrightarrow{T_1 \cup t}$ of $T_1 \cup t$. If $S'_2 - T'_2 \subseteq F$, then the element of $F - (S'_2 - T'_2)$ is loose in Φ ; a contradiction. Thus, at most one element of $S'_2 - T'_2$ is in F , so we may assume that the first two elements of $\overrightarrow{T_1 \cup t}$ are in $(S_1 \cup s) \cap (T_1 \cup t)$. It follows that the first element of $S'_2 - T'_2$ in $\overrightarrow{T_1 \cup t}$ is in the closure or coclosure of $(S_1 \cup s) \cap (T_1 \cup t)$ in $M \setminus x$ and so is loose in Φ ; a contradiction. We conclude that $T_1 \cup t$ is not sequential. A symmetric argument using $T'_2 - S'_2$ and $S'_2 \cap T'_2$ in place of $S'_2 - T'_2$ and $(S_1 \cup s) \cap (T_1 \cup t)$, respectively, establishes that T'_2 is not sequential. Thus $(T'_2, T_1 \cup t)$ is non-sequential; a contradiction. \square

Lemma 5.14. *If $T'_2 \subseteq S'_2$, then $((S_1 \cup s) - t, S_2)$ is a $(4, 4, S)$ -violator for M/t with x in S_2 and $\{s, u, y\} \subseteq (S_1 \cup s) - t$.*

Proof. We have

$$r_{M/s}(S_2) + r_{M/s}(S_1) = r(M/s) + 2.$$

Thus

$$r(S_2 \cup s) - 1 + r(S_1 \cup s) - 1 = r(M/t) + 2.$$

Now, by Lemma 5.4, $r(S_2 \cup s) = r(S_2)$ and $r(T_2 \cup t) = r(T_2)$. Thus, as $T_2 \subseteq S_2$, we deduce that $r(S_2 \cup t) = r(S_2) = r(S_2 \cup s)$, so

$$r(S_2 \cup t) - 1 + r([(S_1 \cup s) - t] \cup t) - 1 = r(M/t) + 2.$$

Hence $((S_1 \cup s) - t, S_2)$ is a 3-separation of M/t .

Evidently $x \in S_2$ and $\{s, u, y\} \subseteq (S_1 \cup s) - t$. Suppose that $((S_1 \cup s) - t, S_2)$ is not a $(4, 4, S)$ -violation of M/t . As (S_1, S_2) is a $(4, 4, S)$ -violation of M/s , it follows that S_1 or S_2 is a quad of M/s . But if S_1 is a quad of M/s , then $S_1 \cup s$ is a circuit of M that properly contains the circuit $\{s, t, u, y\}$; a contradiction. Thus S_2 is a quad of M/s . Hence $S'_2 = T'_2$ since $|S'_2|, |T'_2| \geq 3$, so $S_2 = T_2$ and $((S_1 \cup s) - t, S_2) = (T_1, T_2)$. Thus $((S_1 \cup s) - t, S_2)$ is a $(4, 4, S)$ -violation of M/t . \square

By the last lemma, if $T'_2 \subseteq S'_2$, then we may replace (T_1, T_2) by $((S_1 \cup s) - t, S_2)$ giving $T'_2 = S'_2$. By repeating this process, we may assume that none of S'_2, T'_2 , and U'_2 is properly contained in another such set.

Lemma 5.15. *The sets S'_2, T'_2 , and U'_2 are not all equal.*

Proof. Assume that $S'_2 = T'_2 = U'_2$. We know that $x \in \text{cl}_{M/s}(S'_2) \cap \text{cl}_{M/t}(T'_2) \cap \text{cl}_{M/u}(U'_2)$ and

$$\begin{aligned} \cap(S'_2, \{s, t, u, y\}) &= r(S'_2) + r(\{s, t, u, y\}) - r(S'_2 \cup \{s, t, u, y\}) \\ &= r(S'_2) + 3 - r(S'_2 \cup \{s, t, u, y\}). \end{aligned}$$

Now $M \setminus x$ has $\{s, t, u\}$ as a triad. Thus $r(S'_2 \cup \{s, t, u, y\}) \geq r(S'_2) + 1$. But $\text{cl}(S'_2 \cup s)$ contains x . Thus, by Lemma 5.4 and symmetry, $\text{cl}(S'_2 \cup s)$ contains t and u , and hence y . Therefore $r(S'_2 \cup \{s, t, u, y\}) \leq r(S'_2) + 1$. Thus $\cap(S'_2, \{s, t, u, y\}) = 2$.

By Lemma 2.5,

$$\begin{aligned} \lambda_{M \setminus x}(S'_2 \cup \{s, t, u, y\}) &= \lambda_{M \setminus x}(S'_2) + \lambda_{M \setminus x}(\{s, t, u, y\}) \\ &\quad - \cap_{M \setminus x}(S'_2, \{s, t, u, y\}) - \cap_{M \setminus x}^*(S'_2, \{s, t, u, y\}) \\ &\leq 2 + 2 - 2 = 2. \end{aligned}$$

Since $x \in \text{cl}(S'_2 \cup \{s, t, u, y\})$, we deduce that $\lambda_M(S'_2 \cup \{s, t, u, y, x\}) \leq 2$. As M is 4-connected, it follows that $|E(M) - (S'_2 \cup \{s, t, u, y, x\})| \leq 2$.

By Lemma 2.6(i),

$$2 = \cap(S'_2, S_1 \cup s) \geq \cap(S'_2, \{s, t, u, y\}) = 2.$$

Thus

$$r(S_1 \cup s) - r(S'_2 \cup S_1 \cup s) = r(\{s, t, u, y\}) - r(S'_2 \cup \{s, t, u, y\}).$$

Since $|E(M) - (S'_2 \cup \{s, t, u, y\})| \leq 3$, we deduce that $r(S'_2 \cup S_1 \cup s) = r(S'_2 \cup \{s, t, u, y\}) = r(M)$. Hence $r(S_1 \cup s) = r(\{s, t, u, y\}) = 3$. This contradiction to Lemma 5.3 completes the proof. \square

Lemma 5.16. *If S'_2 and T'_2 are both triads of $M \setminus x$, then $|S'_2 \cap T'_2| = 1$.*

Proof. By Lemma 5.9, $|S'_2 \cap T'_2| \geq 1$. If $|S'_2 \cap T'_2| \geq 2$, then every 3-element subset of $S'_2 \cup T'_2$ is a triad of $M \setminus x$. Thus $r_M^*(S'_2 \cup T'_2) = 3$. Now exactly two elements of D are in $\{s, t, u, y\}$. Thus at most two elements of D are in $S'_2 \cup T'_2$. But, by Lemma 2.9(ii), there is an element of D in each 3-element subset of $S'_2 \cup T'_2$. Hence exactly two elements of D are in $S'_2 \cup T'_2$.

Let $G = S'_2 \cup T'_2 \cup \{s, t, u, y, x\}$. Then G is spanned by $S'_2 \cup T'_2 \cup \{u, x\}$ as $s \in \text{cl}(S_2)$ and $t \in \text{cl}(T_2)$ while $\{s, t, u, y\}$ is a circuit. Thus $r(G) \leq |S'_2 \cup T'_2| = 2$. On the other hand, letting d be an element of $\{s, t, u\}$ such that $|\{d, y\} \cap D| = 1$, we have that $\text{cl}^*(S_2 \cup T_2 \cup \{d, y\})$ contains at least three elements of the cocircuit D and so contains all of D . The choice of D also means that this coclosure contains at least two elements of $\{s, t, u\}$ and the cocircuit $\{s, t, u, x\}$ guarantees that it contains all of $\{s, t, u\}$. Hence $\text{cl}^*(S_2 \cup T_2 \cup \{d, y\})$ contains G and so $r^*(G) \leq r^*(S_2 \cup T_2) + 2 \leq 5$. Thus we have

$$\lambda_M(G) = r(G) + r^*(G) - |G| \leq |S'_2 \cup T'_2| + 2 + 5 - |S'_2 \cup T'_2| + 5 = 2.$$

Hence $|E(M) - G| \leq 2$. But this contradicts the fact that $|E(M)| \geq 12$. \square

Lemma 5.17. *If $|S'_2 \cap T'_2| \geq 2$, then $\lambda_{M \setminus x}(S'_2 \cup T'_2) = 2$. Moreover, if at least two of $S'_2 \cap T'_2, T'_2 \cap U'_2$, and $U'_2 \cap S'_2$ exceed one, then $\lambda_{M \setminus x}(S'_2 \cup T'_2 \cup U'_2) = 2$.*

Proof. We have $\lambda_{M \setminus x}(S'_2) = \lambda_{M \setminus x}(T'_2) = 2$. Since $M \setminus x$ is 3-connected and each of $S'_2 \cap T'_2$ and $E(M \setminus x) - (S'_2 \cup T'_2)$ has at least two elements, the first assertion of the lemma holds by uncrossing.

Now assume that $|S'_2 \cap T'_2| \geq 2$ and $|T'_2 \cap U'_2| \geq 2$. Then $\lambda_{M \setminus x}(S'_2 \cup T'_2) = 2 = \lambda_{M \setminus x}(T'_2 \cup U'_2)$. Since $E(M \setminus x) - (S'_2 \cup T'_2 \cup U'_2) \supseteq \{s, t, u, y\}$, another application of uncrossing gives that $\lambda_{M \setminus x}(S'_2 \cup T'_2 \cup U'_2) = 2$. \square

Lemma 5.18. *If $\lambda_{M \setminus x}(S'_2 \cup T'_2 \cup U'_2) = 2$, then $|(S'_2 \cup T'_2) - U'_2| \geq 2$.*

Proof. Assume that $|(S'_2 \cup T'_2) - U'_2| \leq 1$. By Lemma 5.6, $S'_2 \cup T'_2 \cup U'_2 = E(M) - \{s, t, u, y, x\}$. Hence $|U_1| \leq |\{s, t, y\}| + |(S'_2 \cup T'_2) - U'_2| \leq 3 + 1 = 4$; a contradiction to Lemma 5.3. \square

Lemma 5.19. *If $|S'_2 \cap T'_2| = 1$, then $|S'_2 \cap U'_2| = 1$ and $|T'_2 \cap U'_2| = 1$.*

Proof. By Lemma 5.11, S'_2 and T'_2 are both triads of $M \setminus x$. If $|S'_2 \cap U'_2| = 1$, then U'_2 is also a triad. Hence, by Lemma 5.16, $|T'_2 \cap U'_2| = 1$. Thus we may assume that $|S'_2 \cap U'_2| \geq 2$ and $|T'_2 \cap U'_2| \geq 2$. By Lemma 5.17, $\lambda_{M \setminus x}(S'_2 \cup T'_2 \cup U'_2) = 2$. Then, by Lemma 5.18, $|(S'_2 \cup T'_2) - U'_2| \geq 2$. Since $|S'_2 - U'_2| \leq 1$ and $|T'_2 - U'_2| = 1$, we deduce that $S'_2 - U'_2$ and $T'_2 - U'_2$ are disjoint one-element sets. By Lemma 5.6, $E(M)$ is the disjoint union of the sets $U'_2, S'_2 - U'_2, T'_2 - U'_2$, and $\{s, t, u, y, x\}$.

Since both $(U_1 \cup u, U'_2)$ and $(\{s, t, u, y\}, E(M \setminus x) - \{s, t, u, y\})$ are 3-separations of $M \setminus x$ and $\{s, t, u, y\} \subseteq U_1 \cup u$, we have, by Lemma 2.9(ii) that $|D \cap \{s, t, u, y\}| = 2$ and $|D \cap U'_2| = 2$. Furthermore, by Lemma 2.9(iii), since D meets every triad of $M \setminus x$, we must have $D \cap S'_2 \cap U'_2 \neq \emptyset \neq D \cap T'_2 \cap U'_2$.

Now let $G = S'_2 \cup T'_2 \cup \{s, t, u, y, x\}$ and $R = E(M) - G = U'_2 - (S'_2 \cup T'_2)$. The set G is spanned by $S'_2 \cup T'_2 \cup \{x, u\}$ because $\text{cl}(S_2)$ and $\text{cl}(T_2)$ contain s and t , respectively, and $\text{cl}(\{s, t, u\})$ contains y . Thus $r(G) \leq 7$.

Next we compare $r(R)$ and $r(M)$. In $M|U'_2$, each of $S'_2 \cap U'_2$ and $T'_2 \cap U'_2$ is a union of cocircuits. Thus $r(U'_2) \geq r(R) + 2$. Recall that $(U_1 \cup u, U'_2)$ is a 3-separation of $M \setminus x$. Let $S'_2 - U'_2 = \{s'_2\}$ and $T'_2 - U'_2 = \{t'_2\}$. Since

$s'_2 \in \text{cl}_{M \setminus x}^*(U'_2)$, we have $s'_2 \notin \text{cl}_{M \setminus x}(U'_2)$. Thus $r(U'_2 \cup s'_2) = r(U'_2) + 1$ and $((U_1 \cup u) - s'_2, U'_2 \cup s'_2)$ is a 3-separation of $M \setminus x$. Since $t'_2 \in \text{cl}_{M \setminus x}^*(U'_2 \cup s'_2)$, we have $t'_2 \notin \text{cl}_{M \setminus x}(U'_2 \cup s'_2)$. Thus $r(U'_2 \cup s'_2 \cup t'_2) = r(U'_2) + 2$. Hence $r(U'_2 \cup S'_2 \cup T'_2) \geq r(R) + 4$. The set $\{s, t, u, x\}$ is a cocircuit of M avoiding $U'_2 \cup S'_2 \cup T'_2$. Hence $r(M) \geq r(R) + 5$. As $r(G) \leq 7$ and $R = E(M) - G$, we have

$$\lambda_M(G) \leq 7 + [r(R) - r(M)] \leq 7 - 5 = 2,$$

so $|R| \leq 2$. Thus we get a contradiction since $|G| = 10$ and $|E(M)| \geq 13$. \square

Lemma 5.20. *The set $\{s, t, u, y\}$ is a flat of $M \setminus x$.*

Proof. Assume that $e \in E(M \setminus x) - \{s, t, u, y\}$ and $e \in \text{cl}(\{s, t, u, y\})$. Then $M|\{s, t, u, y, e\} \cong U_{3,5}$. The quad D of M/x contains exactly two elements of $\{s, t, u, y\}$ and exactly two elements of $E(M \setminus x) - \{s, t, u, y, e\}$. Thus $\{s, t, u, y, e\}$ contains a 4-circuit having exactly one element in common with the cocircuit D of M ; a contradiction. \square

Lemma 5.21. $|S'_2 \cap T'_2| \neq 1$.

Proof. Assume the contrary. Then, by Lemma 5.19, $|S'_2 \cap U'_2| = 1 = |T'_2 \cap U'_2|$. By Lemma 5.11, each of S'_2, T'_2 , and U'_2 is a triad of $M \setminus x$. Thus each of S'_2, T'_2 , and U'_2 has exactly four elements, so these sets are quads of $M/s, M/t$, and M/u , respectively. Hence $S_2 \cup s, T_2 \cup t$, and $U_2 \cup u$ are circuits of M . Now D contains exactly two elements of the 3-separating set $\{s, t, u, y\}$ of $M \setminus x$. Hence, without loss of generality, $s \notin D$. Moreover, D meets each of S'_2, T'_2 , and U'_2 . Since D is a cocircuit of M and $S_2 \cup s$ is a circuit of M and these sets meet, it follows that $|D \cap S'_2| = 2$. Thus if $S'_2 \cap T'_2 \cap U'_2 = \emptyset$, then $D \supseteq \{s_t, s_u\}$ where $S'_2 \cap T'_2 = \{s_t\}$ and $S'_2 \cap U'_2 = \{s_u\}$; and if $S'_2 \cap T'_2 \cap U'_2 = \{z\}$, then $D \supseteq \{z, s_2\}$ for some s_2 in $S'_2 - z$.

Let $G = S_2 \cup T_2 \cup U_2 \cup \{s, t, u, y\}$. If $S'_2 \cap T'_2 \cap U'_2 = \emptyset$, let $B_G = \{s, t, u, x\} \cup (T'_2 - S'_2)$; and if $S'_2 \cap T'_2 \cap U'_2 = \{z\}$, let $B_G = \{s, t, u, x\} \cup \{z, t_2, u_2\}$ where $t_2 \in T'_2 - z$ and $u_2 \in U'_2 - z$. Then by using, in order, the circuits $\{s, t, u, y\}, T_2 \cup t, D \cup x, S_2 \cup s$, and $U_2 \cup u$, we get that B_G spans G . Thus $r(G) - |G| \leq -5$.

Now if $S'_2 \cap T'_2 \cap U'_2 = \emptyset$, let $B_G^* = \{s, t, u, y\} \cup \{s_t, t_u\}$ where $\{t_u\} = T'_2 \cap U'_2$; and if $S'_2 \cap T'_2 \cap U'_2 = \{z\}$, let $B_G^* = \{s, t, u, y\} \cup \{z, t_2, u_2\}$. Then by using, in order, the cocircuits $\{s, t, u, x\}, D, S_2, T_2$, and U_2 , we get that B_G^* spans G . Thus if $S'_2 \cap T'_2 \cap U'_2 = \emptyset$, then $r^*(G) \leq 6$, so $\lambda_M(G) \leq 1$ and we get a contradiction since $|E(M) - G| \geq 2$ because $|E(M)| \geq 13$.

If $S'_2 \cap T'_2 \cap U'_2 = \{z\}$, then $r^*(G) \leq 7$, so $\lambda_M(G) \leq 2$. Thus we get a contradiction provided $|E(M) - G| \geq 3$, that is, provided $|E(M)| \geq 15$. But we are only guaranteed that $|E(M)| \geq 13$. We shall now more closely examine the situation in which $S'_2 \cap T'_2 \cap U'_2 = \{z\}$ and show that, in that case too, we will get a contradiction, this time only requiring that $|E(M)| \geq 11$.

For the first time in the proof of this theorem, we consider the element c from the hypothesis such that $\{s, t, u, y, c\}$ is 3-separating in $M \setminus x$. By Lemma 5.20, $c \notin \text{cl}_{M \setminus x}(\{s, t, u, y\})$. Thus $\{s, t, u, y, c\}$ contains a cocircuit

C^* of $M \setminus x$ containing c . Since D contains exactly two elements of $\{s, t, u, y\}$ and exactly two elements of $E(M \setminus x) - \{s, t, u, y, c\}$, we deduce that $c \notin D$. Thus, as $z \in D$, we have

$$c \neq z.$$

Now either C^* or $C^* \cup x$ is a cocircuit of M .

5.21.1. $C^* \cup x$ is a cocircuit of M .

Assume not. Then C^* is a cocircuit of M . But both $\{s, t, u\}$ and C^* are cocircuits of $M \setminus x$, so C^* contains at most two elements of $\{s, t, u\}$. Since $C^* \subseteq \{s, t, u, y, c\}$ and $|C^*| \geq 4$, we deduce that C^* contains exactly two of s, t , and u . Thus C^* meets two of the circuits $S_2 \cup s, T_2 \cup t$, and $U_2 \cup u$ of M . But C^* does not contain z or x and the only element of C^* that can be in $S'_2 \cup T'_2 \cup U'_2$ is c . Since $(S'_2 - z) \cup s, (T'_2 - z) \cup t$, and $(U'_2 - z) \cup u$ are disjoint, we have a contradiction. Hence (5.21.1) holds.

Now the cocircuit $C^* \cup x$ meets each of the circuits $S_2 \cup s, T_2 \cup t$, and $U_2 \cup u$, so C^* meets each of $(S'_2 - z) \cup s, (T'_2 - z) \cup t$, and $(U'_2 - z) \cup u$. But $C^* - c$ avoids $S'_2 \cup T'_2 \cup U'_2$ and C^* does not contain all of s, t , and u . Thus C^* must contain exactly two of s, t , and u . Moreover, for the element w of $\{s, t, u\}$ that is not in C^* , we have $c \in W'_2 - z$.

5.21.2. $y \in C^*$.

Suppose $y \notin C^*$. Then $C^* \cup x = \{s, t, u, y\}$. It follows that $M^* \setminus \{s, t, u, c, x\} \cong U_{3,5}$. Thus, since $|E(M)| \geq 11$, Theorem 1.6 implies that $\{s, t, u, c, x\}$ contains at least two elements e such that $M^* \setminus e$ is internally 4-connected. By assumption, $M^* \setminus x$ is not internally 4-connected. Thus, for some e in $\{s, t, u\}$, the matroid M/e is internally 4-connected. This contradiction to the fact that the theorem fails implies that (5.21.2) holds.

Now we know that C^* contains exactly two of s, t , and u . Moreover, although the symmetry between s, t , and u is broken by the fact that $s \notin D$, we will not use D in the short argument to follow. Thus we may assume that $C^* = \{s, t, y, c\}$ and $c \in U'_2 - z$. Then $\{s, t, y, c, x\}$ and $\{s, t, u, x\}$ are cocircuits of M . Eliminating x , we get that M has a cocircuit D^* containing c and contained in $\{s, t, u, y, c\}$. By orthogonality with the circuits $S_2 \cup s$ and $T_2 \cup t$, we deduce that neither s nor t is in D^* . Thus $|D^*| \leq 3$; a contradiction. \square

On combining Lemmas 5.9, 5.21, 5.17, and 5.6, we immediately get the following.

Lemma 5.22. *Each of $S'_2 \cap T'_2, T'_2 \cap U'_2$, and $S'_2 \cap U'_2$ has at least two elements and is 3-separating. Moreover, $E(M) - (S'_2 \cup T'_2 \cup U'_2) = \{s, t, u, y, x\}$.*

Lemma 5.23. *The sets S'_2 and T'_2 have the following properties.*

- (i) $T'_2 \not\subseteq S'_2$; and
- (ii) $|S'_2 - T'_2| = 1$ or $|T'_2 - S'_2| = 1$.

Proof. Assume that $T'_2 \subseteq S'_2$. Then, by our choice of S'_2, T'_2 , and U'_2 , we have $T'_2 = S'_2$. By Lemmas 5.13 and 5.22 and symmetry, $|U'_2 - T'_2| \leq 1$ or $|T'_2 - U'_2| \leq 1$. If $U'_2 \subseteq T'_2$ or $T'_2 \subseteq U'_2$, then our choice of S'_2, T'_2 , and U'_2 means that $U'_2 = T'_2 = S'_2$, a contradiction to Lemma 5.15. Hence $|U'_2 - T'_2| = 1$ and $|T'_2 - U'_2| = 1$. By Lemma 5.22, $E(M) - (S'_2 \cup T'_2 \cup U'_2) = \{s, t, u, y, x\}$. Thus $|T_1| = 4$, a contradiction to Lemma 5.3. Hence (i) holds. Part (ii) follows immediately from Lemma 5.13. \square




FIGURE 1. The cardinalities $n_s, n_t, n_u, n_{st}, n_{su}$, and n_{tu} .

Now define n_s, n_{st} , and n_{su} to be $|S'_2 - (T'_2 \cup U'_2)|, |(S'_2 \cap T'_2) - U'_2|$, and $|(S'_2 \cap U'_2) - T'_2|$, respectively (see Figure 1). Let n_t, n_u , and n_{tu} be defined similarly.

Lemma 5.24. *After a possible relabelling, either*

- (i) $n_s + n_{su} = n_t + n_{st} = n_u + n_{tu} = 1$; or
- (ii) $n_s + n_{su} = n_u + n_{su} = n_u + n_{tu} = 1$.

Proof. By Lemma 5.23,

$$\begin{aligned} |S'_2 - T'_2| = 1 & \quad \text{or} \quad |T'_2 - S'_2| = 1; \\ |T'_2 - U'_2| = 1 & \quad \text{or} \quad |U'_2 - T'_2| = 1; \text{ and} \\ |U'_2 - S'_2| = 1 & \quad \text{or} \quad |S'_2 - U'_2| = 1. \end{aligned}$$

By symmetry and a possible relabelling, we get that either

- (i) $|S'_2 - T'_2| = |T'_2 - U'_2| = |U'_2 - S'_2| = 1$; or
- (ii) $|S'_2 - T'_2| = |U'_2 - T'_2| = |U'_2 - S'_2| = 1$.

The lemma follows by substitution.

Lemma 5.25. *The following inequalities hold:*

$$\begin{aligned} n_s + n_{st} + n_t &\geq 2; \\ n_t + n_{tu} + n_u &\geq 2; \text{ and} \\ n_u + n_{su} + n_s &\geq 2. \end{aligned}$$

Proof. We have $E(M) - (S'_2 \cup T'_2 \cup U'_2) = \{s, t, u, y, x\}$, so $U_1 = \{s, t, y\} \cup [(S'_2 \cup T'_2) - U'_2]$. As $|U_1| \geq 5$, it follows that $|(S'_2 \cup T'_2) - U'_2| \geq 2$. Hence $n_s + n_{st} + n_t \geq 2$. The second and third inequalities in the lemma follow by symmetry. \square

Lemma 5.26. *At most two of n_s, n_t , and n_u equal one.*

Proof. Suppose that $n_s = n_t = n_u = 1$ and let the elements of $S'_2 - (T'_2 \cup U'_2)$, $T'_2 - (S'_2 \cup U'_2)$, and $U'_2 - (S'_2 \cup T'_2)$ be s' , t' , and u' , respectively. Then both $\{s, t, u, y\}$ and $T'_2 \cup U'_2$ are 3-separating in $M \setminus x$. Hence both $\{s, t, u, y\}$ and $\{s, t, u, y, s'\}$ are 3-separating in $M \setminus x$. Thus, by Lemma 5.20, $s' \in \text{cl}_{M \setminus x}^*(\{s, t, u, y\})$. By symmetry, $\{s', t', u'\} \subseteq \text{cl}_{M \setminus x}^*(\{s, t, u, y\})$. As $(\{s, t, u, y\}, E(M \setminus x) - \{s, t, u, y\})$ is a 3-separation of $(M \setminus x)^*$, we have $r_{(M \setminus x)^*}(\text{cl}_{M \setminus x}^*(\{s, t, u, y\}) \cap (E(M \setminus x) - \{s, t, u, y\})) \leq 2$. Thus $\{s', t', u'\}$ is a triangle in $(M \setminus x)^*$ and hence is a triad in $M \setminus x$. This triad avoids the quad D since D has exactly two elements in each of $S'_2 \cup T'_2 \cup U'_2$, $S'_2 \cup T'_2$, $T'_2 \cup U'_2$, and $S'_2 \cup U'_2$. This contradicts Lemma 2.9(ii). \square

Lemma 5.27. $n_u \neq 1$.

Proof. Suppose $n_u = 1$. Assume first that (i) of Lemma 5.24 holds. Then $n_{tu} = 0$ so, by Lemma 5.25, $n_t = 1$. By the symmetry of (i), we also get $n_s = 1$, so we have a contradiction to Lemma 5.26. Hence we may assume that case (ii) of Lemma 5.24 holds. By that, $n_{su} = 0 = n_{tu}$ and $n_s = 1$. By Lemmas 5.25 and 5.26, $n_t \geq 1$ but $n_t \neq 1$. Hence $n_t \geq 2$, that is, $|T'_2 - (S'_2 \cup U'_2)| \geq 2$. Let s' and u' be the unique elements of $S'_2 - (T'_2 \cup U'_2)$ and $U'_2 - (S'_2 \cup T'_2)$, respectively.

In $M \setminus x$, the set $S'_2 \cup U'_2$ is 3-separating. Hence so is $E - x - (S'_2 \cup U'_2)$. Likewise, T'_2 is 3-separating. The union of T'_2 and $E - x - (S'_2 \cup U'_2)$ avoids $\{s', u'\}$. Hence their intersection $T'_2 - (S'_2 \cup U'_2)$ is 3-separating. Now each of $\{s, t, u, y, s'\}$ and $\{s, t, u, y, u'\}$ is 3-separating in $M \setminus x$ and, by Lemma 5.20, $\{s, t, u, y\}$ is a flat of $M \setminus x$. Thus $\{s', u'\} \subseteq \text{cl}_{M \setminus x}^*(\{s, t, u, y\})$. Hence $\Pi_{M \setminus x}^*(S'_2 \cup U'_2, \{s, t, u, y\}) \geq 2$.

By Lemma 5.4, $x \in \text{cl}(S'_2 \cup s) \cap \text{cl}(U'_2 \cup u)$. By orthogonality with the cocircuit $\{s, t, u, x\}$, we deduce that M has circuits containing $\{x, s\}$ and $\{x, u\}$ that are contained in $S_2 \cup s$ and $U_2 \cup u$. Hence, by circuit elimination, $M \setminus x$ has a circuit contained in $(S'_2 \cup U'_2) \cup \{s, t, u, y\}$ that meets both $S'_2 \cup U'_2$ and $\{s, t, u, y\}$. Thus $\Pi_{M \setminus x}(S'_2 \cup U'_2, \{s, t, u, y\}) \geq 1$. By Lemma 2.5, we get

$$\begin{aligned} 3 &\leq \Pi_{M \setminus x}(S'_2 \cup U'_2, \{s, t, u, y\}) + \Pi_{M \setminus x}^*(S'_2 \cup U'_2, \{s, t, u, y\}) \\ &= \lambda_{M \setminus x}(S'_2 \cup U'_2) + \lambda_{M \setminus x}(\{s, t, u, y\}) - \lambda_{M \setminus x}(S'_2 \cup U'_2 \cup \{s, t, u, y\}) \\ &= 2 + 2 - 2 = 2; \end{aligned}$$

a contradiction. \square

By Lemmas 5.24 and 5.27, $n_u = 0$ and $n_s + n_{su} = 1$. Hence $n_s + n_{su} + n_u = 1$. This contradiction to Lemma 5.25 completes the proof of Theorem 5.1. \square

We are now ready to prove Theorem 3.2 and we begin by restating the result for ease of reference.

Theorem 5.28. *Let M be a 4-connected matroid with $|E(M)| \geq 13$. Then M has an element x such that $M \setminus x$ or M/x is $(4, 4, S)$ -connected.*

Proof. By Theorem 3.3, M has an element x such that $M \setminus x$ or M/x is sequentially 4-connected. By duality, we may assume the former. We may also assume that $M \setminus x$ is not $(4, 4, S)$ -connected so is not weakly 4-connected. Thus, by Lemma 2.9, M/x is weakly 4-connected. Hence M/x is not sequentially 4-connected otherwise the theorem holds.

Because $M \setminus x$ is not weakly 4-connected, it has a 3-separation (X, Y) with $|X|, |Y| \geq 5$. As $M \setminus x$ is sequentially 4-connected, we may assume that X is sequential. Thus we may assume that $|X| = 5$ and X has a sequential ordering $(1, 2, 3, 4, 5)$. Let $Z = \{1, 2, 3, 4\}$. Since M has no triangles, $\{1, 2, 3\}$ is a triad of M .

Suppose first that $4 \in \text{cl}_{M \setminus x}^*(\{1, 2, 3\})$. Then every 3-element subset of Z is a triad of $M \setminus x$. Thus $M^*(Z \cup x) \cong U_{3,5}$. Hence, by Theorem 1.6, for some element z in Z , the matroid $M^* \setminus z$ is internally 4-connected. Hence M/z is internally 4-connected so M/z is $(4, 4, S)$ -connected.

We may now assume that $4 \in \text{cl}_{M \setminus x}(\{1, 2, 3\})$. Then Z is a circuit of M . Consider the 3-separating set $\{1, 2, 3, 4, 5\}$ in $M \setminus x$ and apply Theorem 5.1 taking $(1, 2, 3, 4, 5) = (s, t, u, y, c)$. By that result, for some z in $\{s, t, u\}$, the matroid M/z is $(4, 4, S)$ -connected. This completes the proof of the theorem. \square

Corollary 5.29. *Let M be a 4-connected matroid. Then M has an element x such that $M \setminus x$ or M/x is $(4, 5, S)$ -connected.*

Proof. By Theorem 3.3, M has an element z such that $M \setminus z$ or M/z is sequentially 4-connected. By duality, we may assume the former. If $M \setminus z$ is $(4, 5, S)$ -connected, then the corollary holds. Thus we may assume that $M \setminus z$ is not $(4, 5, S)$ -connected. Hence $M \setminus z$ has a 3-separation (X, Y) with $|X|, |Y| \geq 6$. Thus $|E(M)| \geq 13$. Therefore, by Theorem 5.28, M has an element x such that $M \setminus x$ or M/x is $(4, 4, S)$ -connected and so is $(4, 5, S)$ -connected. \square

6. THE INTERNALLY 4-CONNECTED CASE.

In this section, we establish the main theorem when M is internally 4-connected by proving Theorem 3.6, which, for convenience, is restated below as Theorem 6.3. We begin with an elementary lemma.

Lemma 6.1. *Let M be an internally 4-connected matroid with $|E(M)| \geq 8$.*

- (i) *If e is an element of M that is not in a triad, then $M \setminus e$ is 3-connected.*
- (ii) *Every triad of M avoids every triangle of M .*

Proof. For (i), suppose that $M \setminus e$ has a 2-separation (X, Y) . Then

$$r(X) + r(Y) = r(M \setminus e) + 1$$

and $|X|, |Y| \geq 2$. Since $|E(M)| \geq 8$, we may assume that $|Y| \geq 4$. Then

$$r(X \cup e) + r(Y) \leq r(M \setminus e) + 2.$$

Since M is internally 4-connected, we get a contradiction unless $|X \cup e| = 3 = r(X \cup e)$. In the exceptional case, $X \cup e$ is a triad of M ; a contradiction. Thus (i) holds.

To prove (ii), note that if M has a triad that meets a triangle, then, since $|E(M)| \geq 5$, these sets meet in exactly two elements, so M has a 4-element fan F . But $|F|, |E(M) - F| \geq 4$, so we have a contradiction to the fact that M is internally 4-connected. \square

Next we show that it is a straightforward consequence of earlier results that the main theorem holds for internally 4-connected matroids with at most 12 elements.

Corollary 6.2. *Let M be an internally 4-connected matroid that is not isomorphic to a wheel or whirl of rank three. If $|E(M)| \leq 12$, then M has an element e such that $M \setminus e$ or M/e is $(4, 5, S)$ -connected.*

Proof. The corollary holds by Corollary 5.29 if M is 4-connected. Thus, by duality, we may assume that T has a triangle. Then, by Theorem 3.5, M has an element f such that $M \setminus f$ or M/f is sequentially 4-connected. Since $|E(M)| \leq 12$, it follows that $M \setminus f$ or M/f is $(4, 5, S)$ -connected. \square

Theorem 6.3. *Let M be a $(4, 3, S)$ -connected matroid that is not isomorphic to a wheel or whirl of rank three. Then M has an element e such that $M \setminus e$ or M/e is $(4, 5, S)$ -connected.*

Proof. Assume the theorem fails. Then, by the last result and duality, we may assume that $|E(M)| \geq 13$ and that M has a triangle $\{x, y, z\}$.

By Lemma 6.1, we immediately get the following.

6.3.1. *None of x, y , or z is in a triad of M , and all of $M \setminus x$, $M \setminus y$, and $M \setminus z$ are 3-connected.*

6.3.2. *If $e \in \{x, y, z\}$ and (A, B) is a 3-separation of $M \setminus e$ with $|A| \geq 4$, then $\{x, y, z\} \cap A \neq \emptyset$.*

If $\{x, y, z\} - e \subseteq B$, then $(A, B \cup e)$ is a 3-separation of M in which each side has at least four elements; a contradiction. Thus (6.3.2) holds.

Because the theorem fails, each of $M \setminus x$, $M \setminus y$, and $M \setminus z$ has a $(4, 5, S)$ -violator. For the moment, we shall take (X_1, X_2) , (Y_1, Y_2) , and

(Z_1, Z_2) to be 3-separations of $M \setminus x$, $M \setminus y$, and $M \setminus z$, respectively, with $|X_1|, |X_2|, |Y_1|, |Y_2|, |Z_1|, |Z_2| \geq 4$. Without loss of generality, we shall assume that $y \in X_1$ and $z \in X_2$. We shall also assume that $x \in Y_1 \cap Z_1$. By (6.3.2), $z \in Y_2$ and $y \in Z_2$.

Later we will refine the choices of (X_1, X_2) , (Y_1, Y_2) , and (Z_1, Z_2) , thereby breaking the symmetry between them. At this point, however, we do have symmetry and we will prove various properties of any collection of 3-separations that satisfy the conditions above as well as the additional restrictions imposed by specific lemmas.

Lemma 6.4. *If $(X_1, X_2) \cong (X_1 \cup f, X_2 - f)$ for some element f of X_2 and (X_1, X_2) is a $(4, k, S)$ -violator of $M \setminus x$ with $k \geq 4$, then $(X_1 \cup e, X_2 - e)$ is a $(4, k - 1, S)$ -violator of $M \setminus x$.*

Proof. If (X_1, X_2) is non-sequential, then so is $(X_1 \cup e, X_2 - e)$. If (X_1, X_2) is sequential, then $|X_1|, |X_2| \geq k + 1$, so $|X_1 \cup e|, |X_2 - e| \geq k$. \square

Lemma 6.5. *If (X_1, X_2) is a $(4, 4, S)$ -violator of $M \setminus x$, then $y \in \text{cl}(X_1 - y)$.*

Proof. We have $\lambda_{M \setminus x}(X_1) = 2$. If y is a coloop of $(M \setminus x)|X_1$, then $(X_1 - y, X_2 \cup y) \cong (X_1, X_2)$, so $\lambda_{M \setminus x}(X_1 - y) = 2$. But $X_2 \cup y \supseteq \{y, z\}$, so $\lambda_M(X_1 - y) = 2$. This is a contradiction since, by Lemma 6.4, $(X_1 - y, X_2 \cup y)$ is a $(4, 3, S)$ -violator of $M \setminus x$ with $\{y, z\} \subseteq X_2 \cup y$, so $(X_1 - y, X_2 \cup y \cup x)$ is a $(4, 3, S)$ -violator of M . We deduce that $y \in \text{cl}(X_1 - y)$. \square

Lemma 6.6. *If (Y_1, Y_2) is a $(4, 4, S)$ -violator of $M \setminus y$, then $X_2 \cap Y_1 \neq \emptyset$.*

Proof. Suppose that $X_2 \cap Y_1 = \emptyset$. Then, by Lemma 6.5 and symmetry, $x \in \text{cl}(Y_1 - x)$. But $Y_1 - x \subseteq X_2$, so $x \in \text{cl}(X_1)$; a contradiction. \square

Lemma 6.7. *Let (Y_1, Y_2) be a $(4, 5, S)$ -violator of $M \setminus y$.*

- (i) *If (X_1, X_2) is a $(4, 4, S)$ -violator of $M \setminus x$, then $|X_2 \cap Y_1| \geq 2$.*
- (ii) *If $|X_2 \cap Y_1| = 1$ and $|X_2| = 4$, then $X_2 \cap Y_2$ is a triangle of M and M has a cocircuit containing $(X_2 \cap Y_1) \cup x$ and contained in $X_2 \cup x$.*

Proof. Suppose that $X_2 \cap Y_1 = \{e\}$. If $e \in \text{cl}(X_2 \cap Y_2)$, then $e \in \text{cl}(Y_2)$, so $(Y_1, Y_2) \cong (Y_1 - e, Y_2 \cup e)$. By Lemma 6.4, $(Y_1 - e, Y_2 \cup e)$ is a $(4, 4, S)$ -violator of $M \setminus y$. If x is a coloop of $M|(Y_1 - e)$, then $(Y_1, Y_2) \cong (Y_1 - e - x, Y_2 \cup e \cup x)$ and $(Y_1 - e - x, Y_2 \cup e \cup x)$ is a $(4, 3, S)$ -violator of $M \setminus y$. As $y \in \text{cl}(Y_2 \cup e \cup x)$, we get the contradiction that $(Y_1 - e - x, Y_2 \cup e \cup x \cup y)$ is a $(4, 3, S)$ -violator of M . We deduce that x is not a coloop of $M|(Y_1 - e)$, so $x \in \text{cl}(Y_1 - e - x)$. Hence $x \in \text{cl}(X_1)$, a contradiction.

We may now assume that $e \notin \text{cl}(X_2 \cap Y_2)$, so $e \notin \text{cl}(X_2 - e)$. Hence $(X_1, X_2) \cong (X_1 \cup e, X_2 - e)$ in $M \setminus x$. Now $(X_2 - e) \cap Y_1 = \emptyset$. As $x \in \text{cl}(Y_1 - x)$, we deduce that $x \in \text{cl}(X_1 \cup e)$. Thus $(X_1 \cup e \cup x, X_2 - e)$ is a 3-separation of M . This gives a contradiction provided $|X_2 - e| \geq 4$, that is, provided $|X_2| \geq 5$.

Now suppose that $|X_2| = 4$. Then $X_2 - e = X_2 \cap Y_2$ and this set is a triangle or a triad of M . But $X_2 \cap Y_2$ contains a single element, z , of the

triangle $\{x, y, z\}$. Thus $X_2 \cap Y_2$ is a triangle of M . Hence X_2 is sequential in $M \setminus x$ and so (i) holds. Moreover, $M \setminus x$ has a cocircuit that contains e and is contained in $e \cup (X_2 \cap Y_2)$. Hence M has a cocircuit that contains $\{e, x\}$ and is contained in $\{e, x\} \cup (X_2 \cap Y_2)$. \square

Lemma 6.8. *If $|X_2 \cap Y_1|, |X_1 \cap Y_2| \geq 2$ and $y \in \text{cl}(X_1 - y)$ and $x \in \text{cl}(Y_1 - x)$, then $|X_1 \cap Y_2|, |X_2 \cap Y_1| \in \{2, 3\}$ and $\lambda_M(X_1 \cap Y_2) = 2 = \lambda_M(X_2 \cap Y_1)$. Moreover, if $W \in \{X_1 \cap Y_2, X_2 \cap Y_1\}$ and $|W| = 3$, then W is a triangle or triad of M .*

Proof. We have $2 = \lambda_{M \setminus x}(X_2) \geq \lambda_{M \setminus x, y}(X_2) = \lambda_{M \setminus x, y}(X_1 - y)$ and $2 = \lambda_{M \setminus y}(Y_2) \geq \lambda_{M \setminus x, y}(Y_2) = \lambda_{M \setminus x, y}(Y_1 - x)$. By submodularity,

$$(1) \quad 2+2 \geq \lambda_{M \setminus x, y}(X_2) + \lambda_{M \setminus x, y}(Y_1 - x) \geq \lambda_{M \setminus x, y}(X_2 \cap Y_1) + \lambda_{M \setminus x, y}(X_1 \cap Y_2).$$

Since $z \in X_2 \cap Y_2$ while $y \in \text{cl}(X_1 - y)$ and $x \in \text{cl}(Y_1 - x)$, we have that $\lambda_{M \setminus x, y}(X_2 \cap Y_1) = \lambda_M(X_2 \cap Y_1)$ and $\lambda_{M \setminus x, y}(X_1 \cap Y_2) = \lambda_M(X_1 \cap Y_2)$. As $|X_1 \cap Y_2|, |X_2 \cap Y_1| \geq 2$, we deduce, using (1), that $\lambda_M(X_2 \cap Y_1) = 2$ and $\lambda_M(X_1 \cap Y_2) = 2$. Since M is internally 4-connected, we conclude that each of $X_2 \cap Y_1$ and $X_1 \cap Y_2$ has exactly two or exactly three elements. Moreover, each such set with exactly three elements is a triangle or a triad of M . \square

Lemma 6.9. (i) *Let (X_1, X_2) and (Y_1, Y_2) be $(4, 4, S)$ -violators of $M \setminus x$ and $M \setminus y$, respectively. If $|X_2 \cap Y_1|, |X_1 \cap Y_2| \geq 2$, then $|X_1 \cap Y_2|, |X_2 \cap Y_1| \in \{2, 3\}$ and $\lambda_M(X_1 \cap Y_2) = 2 = \lambda_M(X_2 \cap Y_1)$.*
(ii) *Let (X_1, X_2) and (Y_1, Y_2) be $(4, 5, S)$ -violators of $M \setminus x$ and $M \setminus y$, respectively. Then $|X_1 \cap Y_2|, |X_2 \cap Y_1| \in \{2, 3\}$ and $\lambda_M(X_1 \cap Y_2) = 2 = \lambda_M(X_2 \cap Y_1)$.*

Proof. Let (X_1, X_2) and (Y_1, Y_2) be $(4, 4, S)$ -violators of $M \setminus x$ and $M \setminus y$. Then, by Lemma 6.5 and symmetry, $y \in \text{cl}(X_1 - y)$ and $x \in \text{cl}(Y_1 - x)$. Part (i) follows immediately from Lemma 6.8.

Now let (X_1, X_2) and (Y_1, Y_2) be $(4, 5, S)$ -violators of $M \setminus x$ and $M \setminus y$. By Lemma 6.7(i) and symmetry, $|X_1 \cap Y_2|, |X_2 \cap Y_1| \geq 2$. Part (ii) now follows from part (i). \square

Lemma 6.10. *If (X_1, X_2) is a $(4, 4, S)$ -violator of $M \setminus x$, then $X_2 \cap Y_2 \supsetneq \{z\}$.*

Proof. Suppose that $X_2 \cap Y_2 = \{z\}$. Then, by Lemma 6.5 and symmetry, $z \in \text{cl}(X_2 - z)$. But $X_2 - z \subseteq Y_1$, so $z \in \text{cl}(Y_1)$. Since $x \in Y_1$, we deduce that $y \in \text{cl}(Y_1)$; a contradiction. \square

To this point, we have symmetry between (X_1, X_2) , (Y_1, Y_2) , and (Z_1, Z_2) and this symmetry will be heavily exploited in the argument below as we apply the lemmas we have already proved. We shall now specialize the choices of (X_1, X_2) , (Y_1, Y_2) , and (Z_1, Z_2) . In particular, by Theorem 3.5, since $\{x, y, z\}$ is a triangle of M and M is internally 4-connected having at least 13 elements, we may assume that $M \setminus x$ is sequentially 4-connected. We will take the 3-separation (X_1, X_2) of $M \setminus x$ to have the property that X_2 is sequential and $|X_2| = 6$. Hence (X_1, X_2) is a $(4, 5, S)$ -violator of

$M \setminus x$. We also take the 3-separations (Y_1, Y_2) and (Z_1, Z_2) so that they are $(4, 5, S)$ -violators of $M \setminus y$ and $M \setminus z$, respectively.

Now we want to exploit the symmetry between (X_1, y) and (X_2, z) . Although we have made some special assumptions about X_2 , we do still have symmetry between $(z, X_2, x, X_1, Y_1, y, Y_2)$ and $(y, X_1, x, X_2, Z_1, z, Z_2)$ with respect to the hypotheses of Lemma 6.9. This is easy to see using a Venn diagram. Hence an immediate consequence of Lemmas 6.8 and 6.9 is the following.

Corollary 6.11. $|X_1 \cap Z_1|, |X_2 \cap Z_2| \in \{2, 3\}$ and $\lambda_M(X_1 \cap Z_1) = 2 = \lambda_M(X_2 \cap Z_2)$. Moreover, if $W \in \{X_1 \cap Z_1, X_2 \cap Z_2\}$ and $|W| = 3$, then W is a triangle or triad of M .

Although it will not be needed, it is worth noting at this point that we have the following easy bound on $|E(M)|$, where we recall that M is a counterexample to the theorem.

Lemma 6.12. $|E(M)| \leq 17$.

Proof. We have $|E(M)| = |X_1| + |X_2| + 1 = |X_1| + 7$. Now X_1 is the disjoint union of $X_1 \cap Y_2$, $\{y\}$, $X_1 \cap Y_1 \cap Z_1$, and $X_1 \cap Y_1 \cap Z_2$. By Lemma 6.9, $|X_1 \cap Y_2| \leq 3$ and $|X_1 \cap Y_1 \cap Z_2| \leq |Y_1 \cap Z_2| \leq 3$. Moreover, using Corollary 6.11, we have $|X_1 \cap Y_1 \cap Z_1| \leq |X_1 \cap Z_1| \leq 3$. We conclude that $|X_1| \leq 3 + 1 + 3 + 3 = 10$, so $|E(M)| \leq 17$. \square

To complete the proof of the theorem, we will use the fact that X_2 is sequential. Thus there is a sequential ordering $(1, 2, 3, 4, 5, 6)$ of X_2 . Now, because M is internally 4-connected, we have that $z \in \{1, 2, 3, 4\}$.

Lemma 6.13. *Either*

- (i) $|\{1, 2, 3, 4\} \cap Y_2| = 3$ and $|\{1, 2, 3, 4\} \cap Y_1| = 1$ and $\{1, 2, 3, 4\} \cap Y_2$ is a triangle of M ; or
- (ii) $|\{1, 2, 3, 4\} \cap Y_2| = 2$ and $|\{1, 2, 3, 4\} \cap Y_1| = 2$.

Proof. By Lemma 6.9(ii), since (X_1, X_2) and (Y_1, Y_2) are $(4, 5, S)$ -violators of $M \setminus x$ and $M \setminus y$, respectively, we have $|X_2 \cap Y_1|, |X_1 \cap Y_2| \in \{2, 3\}$.

Now $(X_1 \cup 6, X_2 - 6)$ is a $(4, 4, S)$ -violator of $M \setminus x$. Thus, by Lemma 6.7, $|(X_2 - 6) \cap Y_1| \geq 2$. From the previous paragraph, we have $|(X_1 \cup 6) \cap Y_2| \geq |X_1 \cap Y_2| \geq 2$. Hence, by Lemma 6.8, both $|(X_2 - 6) \cap Y_1|$ and $|(X_1 \cup 6) \cap Y_2|$ are in $\{2, 3\}$. Thus if $6 \in X_2 \cap Y_1$, then $|X_2 \cap Y_1| = 3$ and $|(X_2 - 6) \cap Y_1| = 2$. If $6 \in X_2 \cap Y_2$, then $|X_1 \cap Y_2| = 2$ and, by Lemma 6.10, $|(X_2 - 6) \cap Y_2| \geq 2$.

Consider the position of 5. It is straightforward to see that either

- (a) $|\{1, 2, 3, 4\} \cap Y_2| = 3$ and $|\{1, 2, 3, 4\} \cap Y_1| = 1$; or
- (b) $|\{1, 2, 3, 4\} \cap Y_2| = 2$ and $|\{1, 2, 3, 4\} \cap Y_1| = 2$;

unless $X_2 \cap Y_2 = \{z, 5, 6\}$. Consider the exceptional case. We have $(X_1 \cup \{5, 6\}, X_2 - \{5, 6\})$ as a 3-separation of $M \setminus x$. Now $\lambda_{M \setminus x, y}(Y_1 - x) = 2 = \lambda_{M \setminus x, y}(X_2 - \{5, 6\})$. Thus, by the submodularity of the connectivity function and the positions of x, y , and z , we deduce that $\lambda_{M \setminus x, y}(Y_2 \cap (X_1 \cup \{5, 6\})) =$

$\lambda_M(Y_2 \cap (X_1 \cup \{5, 6\})) = 2$. Since $|Y_2 \cap (X_1 \cup \{5, 6\})| \geq 4$, we have a contradiction to the fact that M is internally 4-connected. We deduce that (a) or (b) holds.

If (a) holds, then, by Lemma 6.7(ii), $\{1, 2, 3, 4\} \cap Y_2$ is a triangle of M . \square

By Lemma 6.9, $|X_2 \cap Y_1|$ is 2 or 3. The rest of the proof considers these two possibilities beginning with the first.

Lemma 6.14. *If $|X_2 \cap Y_1| = 2$, then*

- (i) $6 \in X_2 \cap Y_2$;
- (ii) $|X_1 \cap Y_2| = 2$;
- (iii) $(X_1 \cap Y_2) \cup 6$ is a triangle or a triad of M ; and
- (iv) $(X_2 \cap Y_2) - 6$ is a triangle of M containing z .

Proof. By Lemma 6.13, we have two possibilities for the distribution of the elements of $\{1, 2, 3, 4\}$ in $X_2 \cap Y_1$ and $X_2 \cap Y_2$. Suppose first that $|\{1, 2, 3, 4\} \cap Y_1| = 2$. As $|X_2 \cap Y_2| = 2$, we deduce that $\{5, 6\} \subseteq X_2 \cap Y_2$. Now consider the 3-separation $(X_1 \cup \{5, 6\}, X_2 - \{5, 6\})$ of $M \setminus x$. We have $y \in \text{cl}(X_1 - y)$ and $x \in \text{cl}(Y_1 - x)$. Moreover, $|(X_2 - \{5, 6\}) \cap Y_1| = |X_2 \cap Y_1| \geq 2$ and $|(X_1 \cup \{5, 6\}) \cap Y_2| = |X_1 \cap Y_2| + 2 \geq 4 \geq 2$. Thus, by Lemma 6.8, $|(X_1 \cup \{5, 6\}) \cap Y_2| \in \{2, 3\}$; a contradiction. We conclude that $|\{1, 2, 3, 4\} \cap Y_1| \neq 2$, so $|\{1, 2, 3, 4\} \cap Y_1| = 1$.

By Lemma 6.13, $\{1, 2, 3, 4\} \cap Y_2$ is a triangle of M . We know that $z \in \{1, 2, 3, 4\} \cap Y_2$. Thus z is in a triangle of M contained in $X_2 \cap Y_2$ and avoiding $\{5, 6\}$. We now consider where 5 and 6 are. As $(X_1 \cup 6, X_2 - 6)$ is a $(4, 4, S)$ -violator for $M \setminus x$, we have, by Lemma 6.7, that $|(X_2 - 6) \cap Y_1| \geq 2$. But $|X_2 \cap Y_1| = 2$ by assumption. Thus $6 \in X_2 \cap Y_2$ and $5 \in X_2 \cap Y_1$, so (i) holds. Moreover, $|(X_2 - 6) \cap Y_1| = |X_2 \cap Y_1| = 2$ and $|(X_1 \cup 6) \cap Y_2| = |X_1 \cap Y_2| + 1 \geq 3$. Hence, by Lemma 6.9(i), $|(X_1 \cup 6) \cap Y_2| \in \{2, 3\}$. Thus $|X_1 \cap Y_2| = 2$ and $(X_1 \cup 6) \cap Y_2$ is a triangle or a triad of M , so (ii) and (iii) hold. Part (iv) follows from Lemma 6.13. \square

For the rest of the proof, we shall call the elements of Z_1 *red* and those of Z_2 *green*.

Lemma 6.15. $|X_2 \cap Y_1| = 3$.

Proof. Assume that $|X_2 \cap Y_1| = 2$. From the previous lemma, we may assume that $|X_1 \cap Y_2| = 2$. Let the triangle $(X_2 \cap Y_2) - 6$ be $\{z_1, z_2, z\}$. Since $z \notin \text{cl}(Z_1) \cup \text{cl}(Z_2)$, we may assume that $z_1 \in Z_1$ and $z_2 \in Z_2$. Now, by Lemma 6.14(iii), $(X_1 \cap Y_2) \cup 6$ is a triangle or triad of M . By Lemma 6.9, Y_2 contains two or three red elements. Since z_1 is red, $Y_2 - z_1$ contains either one or two red elements. Thus $(X_1 \cap Y_2) \cup 6$ contains either one green and two red elements, or one red and two green elements. In the first case, we recolour the green element γ of $(X_1 \cap Y_2) \cup 6$ to red. This means replacing (Z_1, Z_2) by $(Z_1 \cup \gamma, Z_2 - \gamma)$, which is a $(4, 4, S)$ -violator of $M \setminus z$. Now $|Y_1 \cap (Z_2 - \gamma)| = |Y_1 \cap Z_2| \geq 2$, while $|(Z_1 \cup \gamma) \cap Y_2| = 4$. This gives a contradiction to Lemma 6.9(i).

We may now assume that $\{y_1, y_2, 6\}$ contains one red and two green elements. In that case, we recolour the red element ρ to green, replacing (Z_1, Z_2) by $(Z_1 - \rho, Z_2 \cup \rho)$, which is a $(4, 4, S)$ -violator. Thus, by Lemma 6.7 and symmetry, $|(Z_1 - \rho) \cap Y_2| \geq 2$; a contradiction to the fact that $|(Z_1 - \rho) \cap Y_2| = 1$. \square

Lemma 6.16. $(X_2 \cap Y_2) - z$ is monochromatic.

Proof. Assume that $(X_2 \cap Y_2) - z$ contains one red and one green element. By Lemma 6.9 and symmetry, X_2 contains either two or three green elements. Thus either

- (i) $X_2 \cap Y_1$ contains one red and two green elements; or
- (ii) $X_2 \cap Y_1$ contains one green and two red elements.

Now $X_2 \cap Y_1$ is a triangle or triad of M .

In case (i), we recolour the red element ρ of $X_2 \cap Y_1$ to green, replacing (Z_1, Z_2) by $(Z_1 - \rho, Z_2 \cup \rho)$. Now $|(Z_2 \cup \rho) \cap X_2| = 4$ and $|(Z_1 - \rho) \cap X_1| \geq 2$, so we get a contradiction to Lemma 6.8.

In case (ii), we recolour the one green element γ of $X_2 \cap Y_1$ to red, replacing (Z_1, Z_2) by the $(4, 4, S)$ -violator $(Z_1 \cup \gamma, Z_2 - \gamma)$. Then, by Lemma 6.7 and symmetry, $|(Z_2 - \gamma) \cap X_2| \geq 2$. But $|(Z_2 - \gamma) \cap X_2| = 1$; a contradiction. \square

Lemma 6.17. $|\{1, 2, 3, 4\} \cap Y_2| = 2$.

Proof. We assume that $|\{1, 2, 3, 4\} \cap Y_2| = 3$. Then $\{1, 2, 3, 4\} \cap Y_2$ is a triangle of M containing z . Since neither Z_1 nor Z_2 spans z , the set $(X_2 \cap Y_2) - z$ must contain one red and one green element; a contradiction to Lemma 6.16. \square

Lemma 6.18. (i) One of 5 and 6 is in $X_2 \cap Y_1$ and the other is in $X_2 \cap Y_2$.

- (ii) $Y_2 \cap (X_1 \cup \{5, 6\})$ is a triangle or triad of M .
- (iii) $|X_1 \cap Y_2| = 2$.

Proof. Part (i) follows immediately from the last lemma and the fact that $|X_2 \cap Y_1| = 3$. Consider the 3-separation $(X_1 \cup \{5, 6\}, X_2 - \{5, 6\})$ of $M \setminus x$. We have $|(X_1 \cup \{5, 6\}) \cap Y_2| = |X_1 \cap Y_2| + 1 \geq 3$ and $|(X_2 - \{5, 6\}) \cap Y_1| = 2$. Also $y \in \text{cl}((X_1 \cup \{5, 6\}) - y)$ and $x \in \text{cl}(Y_1 - x)$. Thus, by Lemma 6.8, $Y_2 \cap (X_1 \cup \{5, 6\})$ is a triangle or triad of M . Moreover, $|X_1 \cap Y_2| = 2$. \square

Lemma 6.19. $|Y_2| = 5$ and (Y_1, Y_2) is non-sequential.

Proof. We have $|Y_2| = |Y_2 \cap X_1| + |Y_2 \cap X_2| = 2 + 3 = 5$. By the choice of (Y_1, Y_2) , we deduce that (Y_1, Y_2) must be non-sequential. \square

Lemma 6.20. The elements of $(X_2 \cap Y_2) - z$ are both red.

Proof. Assume the lemma fails. Then, by Lemma 6.16, both the elements of $(X_2 \cap Y_2) - z$ are green. Now X_2 contains either two or three green elements. Assume the latter. Then, by Lemma 6.8, $X_2 \cap Y_2$ is a triangle or a triad of M . Thus if γ is the green element in $X_2 \cap Y_2$, then $(Y_1 - \gamma, Y_2 \cup \gamma) \cong (Y_1, Y_2)$.

Thus $(Y_1 - \gamma, Y_2 \cup \gamma)$, like (Y_1, Y_2) , is non-sequential, and so is a $(4, 5, S)$ -violator of $M \setminus y$. Hence we could replace (Y_1, Y_2) by $(Y_1 - \gamma, Y_2 \cup \gamma)$. But $|X_2 \cap (Y_1 - \gamma)| = 2$, a contradiction to Lemma 6.15. We conclude that X_2 contains exactly two green elements.

The set Y_2 contains two or three red elements while $|Y_2 \cap X_1| = 2$, so both elements of $Y_2 \cap X_1$ are red. As $(X_1 \cup \{5, 6\}) \cap Y_2$ is a triangle or a triad of M , the element γ' of $\{5, 6\} \cap X_2 \cap Y_2$ can be recoloured red, that is, we replace (Z_1, Z_2) by $(Z_1 \cup \gamma', Z_2 - \gamma')$. Since $|(Z_2 - \gamma') \cap X_2| = |Z_2 \cap X_2| - 1 = 1$, we have a contradiction to Lemma 6.7. \square

Lemma 6.21. *The elements of $X_1 \cap Y_2$ are green.*

Proof. We know that Y_2 contains at most three red elements. Hence $X_1 \cap Y_2$ contains at most one red element. If $X_1 \cap Y_2$ does contain a red element, then, using the triangle or triad $(X_1 \cup \{5, 6\}) \cap Y_2$, we can recolour the other element γ of $X_1 \cap Y_2$ to red, replacing (Z_1, Z_2) by $(Z_1 \cup \gamma, Z_2 - \gamma)$. We now get a contradiction to Lemma 6.8 because $|(Z_1 \cup \gamma) \cap Y_2| = 4$ and $|(Z_2 - \gamma) \cap Y_1| \geq 2$. \square

Since both elements of $X_1 \cap Y_2$ are green, we can recolour the element ρ of $\{5, 6\} \cap X_2$ to green, replacing (Z_1, Z_2) by $(Z_1 - \rho, Z_2 \cup \rho)$. As $|(Z_1 - \rho) \cap Y_2| = 1$, we get a contradiction to Lemma 6.7 that completes the proof of Theorem 6.3. \square

7. FINISHING OFF

This section completes the proof of the main theorem of the paper. Our proof will rely on the following lemma, which is a slight strengthening of a result of Geelen and Whittle [3, Theorem 7.1(i)]. The proof is a minor modification of their proof and is presented here for completeness.

Lemma 7.1. *Let M be a sequentially 4-connected matroid and let (A, B) be a sequential 3-separation of M having (a_1, a_2, \dots, a_k) as a sequential ordering of A with $k = |A| \geq 4$. If $M \setminus a_i$ is 3-connected, then $M \setminus a_i$ is sequentially 4-connected.*

Proof. The proof will make repeated use of the elementary observation that if (J, K) is a 3-separating partition of M and $e \in J$, then $(J - e, K)$ is a 3-separating partition of $M \setminus e$. Assume that $M \setminus a_i$ is not sequentially 4-connected, letting (X, Y) be a non-sequential 3-separation of it. Since the first three elements of (a_1, a_2, \dots, a_k) can be arbitrarily reordered, we may assume that $i \geq 3$. Suppose first that $i = 3$. Then $\{a_1, a_2, a_3\}$ is a triangle, otherwise it is a triad and $M \setminus a_3$ is not 3-connected. If $a_4 \in \text{cl}(\{a_1, a_2, a_3\})$, then we can interchange a_3 and a_4 to reduce to the case when $i \geq 4$, which we treat below. If $a_4 \notin \text{cl}(\{a_1, a_2, a_3\})$, then $a_4 \in \text{cl}^*(\{a_1, a_2, a_3\})$. Thus $\{a_1, a_2, a_3, a_4\}$ contains a cocircuit of M containing a_4 . Since $M \setminus a_3$ is 3-connected, it has $\{a_1, a_2, a_4\}$ as a triad. Now at least two of a_1, a_2 , and a_4 may be assumed to be in X , so $(X \cup \{a_1, a_2, a_4\}, Y - \{a_1, a_2, a_4\})$ is a non-sequential 3-separation of $M \setminus a_3$. Thus $(X \cup \{a_1, a_2, a_3, a_4\}, Y - \{a_1, a_2, a_4\})$

is a 3-separation of M . This 3-separation must be sequential so, by Lemma 2.7, $(X \cup \{a_1, a_2, a_4\}, Y - \{a_1, a_2, a_4\})$ is a sequential 3-separation of $M \setminus a_3$; a contradiction.

Now suppose that $i \geq 4$. We may assume that at least two of a_1, a_2 , and a_3 are in X . Hence each of $X \cup \{a_1, a_2, a_3\}$, $X \cup \{a_1, a_2, a_3, a_4\}$, \dots , $X \cup \{a_1, a_2, \dots, a_{i-1}\}$ is 3-separating in $M \setminus a_i$, so $(X \cup \{a_1, a_2, \dots, a_{i-1}\}, Y - \{a_1, a_2, \dots, a_{i-1}\})$ is a non-sequential 3-separation of $M \setminus a_i$. Now $a_i \in \text{cl}(\{a_1, a_2, \dots, a_{i-1}\})$, or $a_i \in \text{cl}^*(\{a_1, a_2, \dots, a_{i-1}\})$. In the latter case, $r(\{a_1, a_2, \dots, a_i\}) = r(\{a_1, a_2, \dots, a_{i-1}\}) + 1$, so $\lambda_{M \setminus a_i}(\{a_1, a_2, \dots, a_{i-1}\}) = 1$; a contradiction. Therefore $a_i \in \text{cl}(\{a_1, a_2, \dots, a_{i-1}\})$ and $(X \cup \{a_1, a_2, \dots, a_i\}, Y - \{a_1, a_2, \dots, a_{i-1}\})$ is a 3-separation of M . This 3-separation must be sequential, yet this implies, by Lemma 2.7, that $(X \cup \{a_1, a_2, \dots, a_{i-1}\}, Y - \{a_1, a_2, \dots, a_{i-1}\})$ is a sequential 3-separation of $M \setminus a_i$; a contradiction. \square

Next we prove the main theorem in the case that M is $(4, 4)$ -connected.

Theorem 7.2. *Let M be a $(4, 4, S)$ -connected matroid that is not isomorphic to a wheel or whirl of rank 3 or 4. Then M has an element x such that $M \setminus x$ or M/x is $(4, 5, S)$ -connected.*

Proof. By Theorem 6.3, the result holds if M is $(4, 3, S)$ -connected. Thus we may assume that M has a 3-separation (X, Y) with $|X| = 4$ and $|Y| \geq 4$ and with X sequential. Let (x_1, x_2, x_3, x_4) be a sequential ordering of X . Then $\{x_1, x_2, x_3\}$ is a triangle or a triad of M . By duality, we may assume that $x_4 \in \text{cl}(\{x_1, x_2, x_3\})$. Then it is straightforward to show that $(\{x_1, x_2, x_3\}, E(M) - \{x_1, x_2, x_3, x_4\})$ is a non-minimal 2-separation of M/x_4 . Hence, by Lemma 2.4, $\text{co}(M \setminus x_4)$ is 3-connected. Thus either

- (i) $M \setminus x_4$ is 3-connected, or
- (ii) M has a triad T^* containing x_4 .

Consider case (ii). As $x_4 \in \text{cl}(\{x_1, x_2, x_3\})$, the triad T^* meets $\{x_1, x_2, x_3\}$. If $T^* \subseteq \{x_1, x_2, x_3, x_4\}$, then $\lambda_M(\{x_1, x_2, x_3, x_4\}) = 1$; a contradiction. Hence $|T^* \cap \{x_1, x_2, x_3, x_4\}| = 2$ so, by Lemma 2.3, $T^* \cup \{x_1, x_2, x_3, x_4\}$ is 3-separating in M . If $|E(M)| \geq 10$, then $|E(M) - (T^* \cup \{x_1, x_2, x_3, x_4\})| \geq 5$, so we have a contradiction to the fact that M is $(4, 4, S)$ -connected. Now assume that $|E(M)| < 10$. We know that $|E(M)| \geq 8$. Hence, by Theorem 1.2, either M is a wheel or whirl of rank 4, or M has an element e such that $M \setminus e$ or M/e is sequentially 4-connected. The former case was excluded by assumption. In the latter case, because $|E(M)| < 13$, either $M \setminus e$ or M/e is $(4, 5, S)$ -connected.

Now consider case (i). By Lemma 7.1, $M \setminus x_4$ is sequentially 4-connected. Suppose this matroid has a 3-separation (J, K) with $|J|, |K| \geq 6$. Without loss of generality, at least two of x_1, x_2 , and x_3 are in J . Thus $(J, K) \cong (J \cup \{x_1, x_2, x_3\}, K - \{x_1, x_2, x_3\})$ and $(J \cup \{x_1, x_2, x_3, x_4\}, K - \{x_1, x_2, x_3\})$ is a 3-separation of M . Since $|J \cup \{x_1, x_2, x_3, x_4\}|, |K - \{x_1, x_2, x_3\}| \geq 5$, we

have a contradiction to the fact that M is $(4, 4, S)$ -connected. We conclude that $M \setminus x_4$ is $(4, 5, S)$ -connected. \square

To complete the proof of the main theorem, we shall require some more preliminaries some of which are extracted from Hall's proof of Theorem 1.3. A *segment* in a matroid N is a subset X of $E(N)$ such that every 3-element subset of X is a circuit of N . A *cosegment* of N is a segment of N^* .

Lemma 7.3. [4, Lemma 4.1] *If M is a $(4, k)$ -connected matroid and X is a 4-element segment, then $M \setminus x$ is $(4, k)$ -connected for some x in X .*

Lemma 7.4. *Let A be a 5-element sequential 3-separating set in a $(4, 5, S)$ -connected matroid M having at least 13 elements. Let $(a_1, a_2, a_3, a_4, a_5)$ be a sequential ordering of A . If $i \in \{1, 2, 3\}$ and $\{a_1, a_2, a_3\}$ is a triangle, or if $i \geq 4$ and $a_i \in \text{cl}(\{a_1, a_2, \dots, a_{i-1}\})$, then $M \setminus a_i$ is 3-connected unless a_i is in a triad of M contained in A .*

Proof. Suppose first that $i \geq 4$ and $a_i \in \text{cl}(\{a_1, a_2, \dots, a_{i-1}\})$. Then M/a_i has $(\{a_1, a_2, \dots, a_{i-1}\}, \{a_{i+1}, a_5\} \cup B)$ as a non-minimal 2-separation. Hence, by Lemma 2.4, $\text{co}(M \setminus a_i)$ is 3-connected. Thus $M \setminus a_i$ is 3-connected unless a_i is in a triad T^* of M . In the exceptional case, as $a_i \in \text{cl}(\{a_1, a_2, \dots, a_{i-1}\})$, it follows by orthogonality that T^* meets $\{a_1, a_2, \dots, a_{i-1}\}$. Thus T^* and A are 3-separating in M having at least two common elements. Therefore $T^* \cup A$ is 3-separating. If $T^* \not\subseteq A$, then $|T^* \cup A| = 6$ and so we contradict the fact that M is $(4, 5, S)$ -connected. Hence, when $i \geq 4$, the matroid $M \setminus a_i$ is 3-connected unless a_i is in a triad of M contained in A .

Now assume that $i \in \{1, 2, 3\}$ and $\{a_1, a_2, a_3\}$ is a triangle. Since a_1, a_2 , and a_3 can be arbitrarily reordered, we may assume that $i = 1$. Suppose that (X, Y) is a non-minimal 2-separation of $M \setminus a_1$. If $a_4 \in \text{cl}(\{a_1, a_2, a_3\})$, then $\{a_1, a_2, a_3, a_4\}$ is a segment so, by Lemma 7.3, $M \setminus a_1$ is 3-connected. We may now assume that $a_4 \in \text{cl}^*(\{a_1, a_2, a_3\})$. Then M has a cocircuit C^* containing a_4 and contained in $\{a_1, a_2, a_3, a_4\}$. Suppose that $|C^*| = 4$. Then $\{a_2, a_3, a_4\}$ is a cocircuit of $M \setminus a_1$. We may assume that at least two elements of $\{a_2, a_3, a_4\}$ are in X . Thus $(X \cup \{a_2, a_3, a_4\}, Y - \{a_2, a_3, a_4\})$ is a 2-separation of $M \setminus a_1$. Hence $(X \cup \{a_1, a_2, a_3, a_4\}, Y - \{a_2, a_3, a_4\})$ is a 2-separation of M . But M is 3-connected, so $|Y - \{a_2, a_3, a_4\}| < 2$, which contradicts the fact that $|Y| \geq 3$. We conclude that the only 2-separations of $M \setminus a_1$ are minimal. Hence either $M \setminus a_i$ is 3-connected, or a_1 is in a triad T^* of M . In the latter case, we argue as at the end of the previous paragraph to deduce that $T^* \subseteq A$. \square

The next lemma and its proof are lifted from Hall [4, p. 56].

Lemma 7.5. *Let M be a $(4, 5)$ -connected matroid with $|E(M)| \geq 16$. Let A be a 5-element 3-separating set in M with $r(A) = 3$. If a is an element of A for which $M \setminus a$ is 3-connected and $A - a$ contains no triangles, then $M \setminus a$ is $(4, 5)$ -connected.*

Proof. Assume that $M \setminus a$ has a 3-separation (X, Y) with $|X|, |Y| \geq 6$. Since $A - a$ contains no triangles and $r(A) = 3$, every 3-element subset of $A - a$ spans A . Since neither $\text{cl}(X)$ nor $\text{cl}(Y)$ contains a , we deduce that $|A \cap X| = 2 = |A \cap Y|$. Since $M \setminus a$ is 3-connected, $\lambda_{M \setminus a}(A \cap X) = 2 = \lambda_{M \setminus a}(A \cap Y)$. Thus, by the submodularity of λ , we deduce that both $Y \cap (E(M) - A)$ and $X \cap (E(M) - A)$ are 3-separating in $M \setminus a$. Because $a \in \text{cl}(A - a)$, these sets are also 3-separating in M . Thus $|X \cap (E(M) - A)|, |Y \cap (E(M) - A)| \leq 5$. Since $|A| = 5$, it follows that $|E(M)| \leq 15$; a contradiction. We conclude that $M \setminus a$ is $(4, 5)$ -connected. \square

Lemma 7.6. *Let M be a $(4, 5, S)$ -connected matroid with $|E(M)| \geq 12$. Let $\{a_1, a_2, a_3, a_4, a_5\}$ be a 5-element fan F in M having $\{a_1, a_2, a_3\}$ and $\{a_3, a_4, a_5\}$ as triangles and $\{a_2, a_3, a_4\}$ as a triad. Then $M \setminus a_3/a_4$ is sequentially 4-connected.*

Proof. If a_4 is in a triangle T other than $\{a_3, a_4, a_5\}$, then, by orthogonality and the fact that M is 3-connected, it follows that $T = \{a_2, a_4, a_6\}$ for some new element a_6 . Then $F \cup T$ is a 6-element 3-separating set in M ; a contradiction since $|E(M)| \geq 12$. We deduce that $\{a_3, a_4, a_5\}$ is the unique triangle containing a_4 . A similar argument (or see [6, Lemma 3.4]) establishes that $\{a_2, a_3, a_4\}$ is the unique triad of M containing a_3 . Hence if $M \setminus a_3/a_4$ is not 3-connected, it has a 2-separation (J, K) with $|J|, |K| \geq 3$. On the other hand, if $M \setminus a_3/a_4$ is 3-connected but not sequentially 4-connected, it has a non-sequential 3-separation (J, K) . We shall prove simultaneously that $M \setminus a_3/a_4$ is 3-connected and that it is sequentially 4-connected by considering a k -separation (J, K) of $M \setminus a_3/a_4$ for some $k \in \{2, 3\}$, where $|J|, |K| \geq 3$ if $k = 2$, while (J, K) is non-sequential if $k = 3$.

We may assume that at least two elements of $\{a_1, a_2, a_5\}$ are in J , so $(J \cup \{a_1, a_2, a_5\}, K - \{a_1, a_2, a_5\})$ is a k -separation of $M \setminus a_3/a_4$. Moreover, if $k = 3$, this 3-separation is non-sequential while if $k = 2$, then $|K - \{a_1, a_2, a_5\}| \geq 2$. Hence $(J \cup \{a_1, a_2, a_5, a_3\}, K - \{a_1, a_2, a_5\})$ is a k -separation of M/a_4 . As $a_4 \in \text{cl}^*(\{a_2, a_3\})$, it follows that $(J \cup F, K - F)$ is a k -separation of M . If $k = 2$, then, as $|K - F| \geq 2$, we contradict the fact that M is 3-connected. We conclude $M \setminus a_3/a_4$ is 3-connected. If $k = 3$, then, since M is sequentially 4-connected, $(J \cup F, K - F)$ is a sequential 3-separation of M . Thus, by Lemma 2.7, $(J \cup \{a_1, a_2, a_5\}, K - \{a_1, a_2, a_5\})$ is a sequential 3-separation of $M \setminus a_3/a_4$; a contradiction. We conclude that $M \setminus a_3/a_4$ is sequentially 4-connected. \square

We are now ready to complete the proof of the main theorem of the paper.

Proof of Theorem 3.1. If M is $(4, 4, S)$ -connected, then the theorem follows by Theorem 7.2. We may now assume that M is $(4, 5, S)$ -connected but not $(4, 4, S)$ -connected. Then M has a 3-separation (A, B) with $|A|, |B| \geq 5$. Since M is sequentially 4-connected, we may assume that A is sequential having exactly 5 elements.

Suppose that A contains a 4-element segment. Then, by Lemma 7.3, A contains an element e such that $M \setminus e$ is $(4, 5)$ -connected. In particular, $M \setminus e$ is 3-connected so, by Lemma 7.1, $M \setminus e$ is sequentially 4-connected. Hence $M \setminus e$ is $(4, 5, S)$ -connected and the theorem holds.

By the last paragraph and duality, we may assume that A contains no 4-element segments or cosegments of M . By Theorem 1.2, either M is neither a wheel nor a whirl and M has an element e such that $M \setminus e$ or M/e is sequentially 4-connected; or M is a wheel or a whirl and $\text{co}(M \setminus e)$ or $\text{si}(M/e)$ is sequentially 4-connected for every element e . Since a sequentially 4-connected matroid N is certainly $(4, 5, S)$ -connected when $|E(N)| \leq 12$, we deduce that the theorem holds when $|E(M)| \leq 12$. Thus we may assume that $|E(M)| \geq 13$. Hall's proof of Theorem 1.3 distinguishes the cases when $|E(M)| \geq 16$ and when $13 \leq |E(M)| \leq 15$, and, since we will be relying on her results, we shall use the same dichotomy.




FIGURE 2. The five possibilities for the 3-separating set A .

Since A is a 3-separating set in M , we have $r(A) + r^*(A) - |A| = 2$, so $r(A) + r^*(A) = 7$. Because A contains no 4-element segments or cosegments of M , we may assume by duality that $r(A) = 3$. Moreover, since M is $(4, 5)$ -connected, A is a flat of M . Hall [4, p. 58] distinguishes eleven possibilities for A . Since we have the additional requirement that A is sequential, we can reduce the number of possibilities to five. In particular, using Hall's terminology, A is a 5-element fan or a 3-separating set of type A, type B, type D, or type F. In each case, we have labelled A in Figure 2 such that $(a_1, a_2, a_3, a_4, a_5)$ is a sequential ordering of it. To interpret this

diagram, observe that, in each case, we have drawn $M|A$. The line in the diagram marking the boundary between the plane A and the hyperplane $\text{cl}(B)$ corresponds to $\text{cl}(A) \cap \text{cl}(B)$.

Suppose that $|E(M)| \geq 16$. If A is a 5-element fan, then, by Hall [4, pp. 57–58], either $M \setminus a_1$ or $M \setminus a_3/a_4$, which is isomorphic to $\text{co}(M \setminus a_3)$, is $(4, 5)$ -connected. Hence, by Lemmas 7.1 and 7.6, $M \setminus a_1$ or $\text{co}(M \setminus a_3)$ is $(4, 5, S)$ -connected. We may now assume that A has type A, B, D, or F. By Lemma 7.4, taking $i = 4$ when A has type A or B and taking $i = 3$ when A has type D or F, we see that the matroid $M \setminus a_i$ is 3-connected. Thus, by Lemma 7.1 $M \setminus a_i$ is sequentially 4-connected. Moreover, by Lemma 7.5, $M \setminus a_i$ is $(4, 5)$ -connected and so is $(4, 5, S)$ -connected. We conclude that the theorem holds when $|E(M)| \geq 16$.

Now suppose that $13 \leq |E(M)| \leq 15$. In this case, if A is a fan, then, by Hall [4, 5.2.10], one of $M \setminus a_1$, $M \setminus a_5$, or $\text{co}(M \setminus a_3)$ is $(4, 5)$ -connected. Again, by Lemmas 7.1 and 7.6, $M \setminus a_1$, $M \setminus a_5$, or $\text{co}(M \setminus a_3)$ is $(4, 5, S)$ -connected. Now assume that A has type A, B, D, or F. In each of these cases, Hall identified a pair of elements $\{a_i, a_j\}$ such that $M \setminus a_i$ or $M \setminus a_j$ is $(4, 5)$ -connected. In particular, $\{i, j\}$ is $\{4, 5\}$ if A has type A or B [4, 5.2.2, 5.2.3]; $\{i, j\}$ is $\{2, 3\}$ if A has type D [4, 5.2.4]; and $\{i, j\}$ is $\{3, 5\}$ if A has type F [4, 5.2.5]. By Lemma 7.1, we get that $M \setminus a_i$ or $M \setminus a_j$ is $(4, 5, S)$ -connected and this completes the proof of the theorem. \square




FIGURE 3. Simplification and cosimplification are needed.

It is natural to ask whether there is a $(4, 5, S)$ -connected matroid M other than a wheel or a whirl in which there is no element e such that $M \setminus e$ or M/e is $(4, 5, S)$ -connected. In other words, are we forced to allow cosimplification or simplification in Theorem 1.5? The cycle matroid M of the graph G in Figure 3 is $(4, 5, S)$ -connected. All 18 elements of M lie in triangles. Nine of the elements, including all those bounding the infinite face F of G , also

lie in triads. The remaining nine elements are of two types: those that meet a degree-4 vertex on the boundary of F ; and those bounding the innermost triangular face of G . The deletion of an edge of the first type creates a 6-element fan, while deletion of an edge of the second type leaves a 3-vertex cut corresponding to a 3-separation in which each part has 8 or 9 elements.

REFERENCES

- [1] Bixby, R. E., A simple theorem on 3-connectivity, *Linear Algebra Appl.* **45** (1982), 123–126.
- [2] Coullard, C. R., *Minors of 3-connected matroids and adjoints of binary matroids*, Ph.D. thesis, Northwestern University, 1985.
- [3] Geelen, J. and Whittle, G., Matroid 4-connectivity: A deletion-contraction theorem, *J. Combin. Theory Ser. B* **83** (2001), 15–37.
- [4] Hall, R., A chain theorem for 4-connected matroids, *J. Combin. Theory Ser. B* **93** (2005), 45–66.
- [5] Oxley, J. G., *Matroid Theory*, Oxford University Press, New York, 1992.
- [6] Oxley, J., and Wu, H., On the structure of 3-connected matroids and graphs, *European J. Combin.* **21** (2000), 667–688.
- [7] Oxley, J., Semple, C., and Whittle, G., The structure of the 3-separations of 3-connected matroids, *J. Combin. Theory Ser. B* **92** (2004), 257–293.
- [8] Oxley, J., Semple, C., and Whittle, G., The structure of the 3-separations of 3-connected matroids II, *European J. Combin.*, to appear.
- [9] Rajan, A., Algorithmic applications of connectivity and related topics in matroid theory, Ph.D. thesis, Northwestern University, 1987.
- [10] Tutte, W. T., Connectivity in matroids, *Canad. J. Math.* **18** (1966), 1301–1324.

DEPARTMENT OF MATHEMATICS, LOUISIANA STATE UNIVERSITY, BATON ROUGE,
LOUISIANA, USA
E-mail address: oxley@math.lsu.edu

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF CANTERBURY,
CHRISTCHURCH, NEW ZEALAND
E-mail address: c.semple@math.canterbury.ac.nz

SCHOOL OF MATHEMATICAL AND COMPUTING SCIENCES, VICTORIA UNIVERSITY,
WELLINGTON, NEW ZEALAND
E-mail address: Geoff.Whittle@mcs.vuw.ac.nz